

MORITA CLASSES OF MICRODIFFERENTIAL ALGEBROIDS

ANDREA D'AGNOLO AND PIETRO POLESELLO

Dedicated to the memory of Louis Boutet de Monvel

ABSTRACT. Projective cotangent bundles of complex manifolds are the local models of complex contact manifolds. Such bundles are quantized by the algebra of microdifferential operators (a localization of the algebra of differential operators on the base manifold).

Kashiwara proved that any complex contact manifold X is quantized by a canonical microdifferential algebroid (a linear stack locally equivalent to an algebra of microdifferential operators). Besides the canonical one, there can be other microdifferential algebroids on X . Our aim is to classify them. More precisely, let Y be the symplectification of X . We prove that Morita (resp. equivalence) classes of microdifferential algebroids on X are described by $H^2(Y; \mathbb{C}^\times)$. We also show that any linear stack locally equivalent to a stack of microdifferential modules is in fact a stack of modules over a microdifferential algebroid.

To obtain these results we use techniques of microlocal calculus, non-abelian cohomology and Morita theory for linear stacks.

CONTENTS

Introduction	2
1. Non-abelian cohomology	4
2. Algebroids	7
3. Morita theory for linear stacks	13
4. Microdifferential operators	23
5. Microdifferential algebroids	29
Appendix A. Cocycles	36
References	40

2010 *Mathematics Subject Classification.* 58J15, 53C56, 32C38.

Key words and phrases. Morita equivalence, algebroid stacks, microdifferential operators, quantization, contact manifolds.

Partial support from the Fondazione Cariparo through the project “Differential methods in Arithmetic, Geometry and Algebra”, from Padova University through the grant CPDA122824/12, and from E.S.I. program “Higher Structures in Mathematics and Physics” are acknowledged. The first author wishes to thank R.I.M.S. of Kyoto University for kind hospitality during the preparation of this paper. The second author wishes to thank E.S.I. of Vienna and Paris 6 University for kind hospitality during the preparation of this paper.

INTRODUCTION

In recent years there has been a lot of interest in the study of the deformation-quantizations of complex/algebraic symplectic manifolds, or more generally Poisson manifolds (see for example [26, 35, 4, 33, 9, 5, 23, 41]). We are interested here in the study of (non-formal) quantizations of complex contact manifolds, which are locally modeled on the algebra of microdifferential operators of [37], as initiated by Kashiwara [17] (we refer to [35, 20, 10] for the relations between such quantizations and deformation-quantizations).

Let us describe such quantizations in a few more details.

Let X be a complex contact manifold. By Darboux theorem, a local model of X is an open subset of the projective cotangent bundle P^*M of a complex manifold M . Let \mathcal{E}_{P^*M} be the sheaf of microdifferential operators on P^*M (a localization of the algebra of differential operators on M). A microdifferential algebra (\mathcal{E} -algebra, for short) on X is a sheaf of \mathbb{C} -algebras locally isomorphic to \mathcal{E}_{P^*M} .

To quantize X in the strict sense means to endow it with an \mathcal{E} -algebra. (The relation between quantization and microdifferential operators is classical, see for example [2, 34] for details.) This might not be possible in general. However, Kashiwara [17] proved that X is endowed with a canonical \mathcal{E} -algebroid \mathbf{E}_X . This means the following. To an algebra A one associates the linear category with one object and elements of A as its endomorphisms. Similarly, to a sheaf of algebras on X one associates a linear stack. An \mathcal{E} -algebroid on X is a \mathbb{C} -linear stack locally equivalent to one associated with an \mathcal{E} -algebra. In this sense, a quantization of X is the datum of an \mathcal{E} -algebroid on X .

Having to deal with an algebroid instead of an algebra is not very restrictive. For example, it makes sense to consider coherent modules over an \mathcal{E} -algebroid, and in particular regular modules along complex involutive subvarieties of X . The Lagrangian case is of particular interest, since these modules are the counterpart of microlocal perverse sheaves in the Riemann-Hilbert correspondence (see [17, 39, 12, 13]). A very interesting example of non-coherent module is the (twisted) sheaf of microfunctions along a totally real, I-symplectic Lagrangian submanifold (see [17, §4]).

A natural problem is to classify \mathcal{E} -algebroids on X .

The canonical \mathcal{E} -algebroid \mathbf{E}_X is endowed with an anti-involution, corresponding to the operation of taking the formal adjoint of microdifferential operators. It is also endowed with a natural order filtration, and its associated graded algebroid is trivial. It is shown in [32] that \mathbf{E}_X is unique among such \mathcal{E} -algebroids.

Equivalence classes of filtered \mathcal{E} -algebroids with trivial graded (but not necessarily endowed with an anti-involution) are classified in [34].

In this paper, we classify arbitrary \mathcal{E} -algebroids on X . These include in particular filtered \mathcal{E} -algebroids whose associated graded is non-trivial (what would be called twisted quantizations of X in the terminology of [4, 41]).

We also classify stacks of twisted \mathcal{E} -modules, i.e. stacks locally equivalent to the stack of modules over an \mathcal{E} -algebra. These are the most general stacks where the notion of microdifferential modules makes sense. Roughly, they are obtained by patching sheaves of \mathcal{E} -algebras through Morita equivalences.

More precisely, in Theorems 5.2.3 and 5.4.3, and Corollary 5.4.2, we prove the following results:

- (i) Two \mathcal{E} -algebroids are equivalent if and only if they are Morita equivalent, i.e. their stacks of modules are equivalent.
- (ii) Any stack of twisted \mathcal{E} -modules is globally equivalent to the stack of modules over an \mathcal{E} -algebroid.
- (iii) The set of equivalence classes (resp. Morita classes) of \mathcal{E} -algebroids is canonically isomorphic to $H^2(Y; \mathbb{C}_Y^\times)$, for Y the symplectification of X .
- (iv) The group of invertible \mathbf{E}_X -bimodules is isomorphic to $H^1(Y; \mathbb{C}_Y^\times)$.

We also give explicit realizations of the isomorphisms in (iii) and (iv).

To obtain our results, we use techniques of microlocal calculus, non-abelian cohomology and Morita theory for linear stacks.

The content of this paper is as follows.

In Section 1 we collect, without proofs, the main facts of non-abelian cohomology we need to prove our results. Note that cohomology with values in non-abelian groups was already used in [2] to classify \mathcal{E} -algebras, and cohomology with values in 2-groups is used in [33, 34] for the classification of algebroids.

In Section 2 we give the basics of the theory of algebroids.

In Section 3, we detail Morita theory for linear stacks. In particular, the notion of Picard good stacks allows us to recover results of [27]. Morita theory for linear categories is developed in [29, 31]. The case of stacks of modules over sheaves of algebras is discussed in [22] (see also [11]).

In Section 4 we recall some results from the theory of microdifferential operators. In particular, we provide a detailed proof of an unpublished result, due to Kashiwara, on the structure of invertible bimodules (see Theorem 4.3.5 below). This allows us to prove the key Theorem 4.3.6.

In Section 5 we prove our main classification results, as (i)–(iv) above.

In Appendix A we recall the cocycle description of algebroids and functors between them.

For symplectic manifolds, or more generally for Poisson manifolds, some results related to ours appeared in the literature: on a complex symplectic manifold, deformation quantization algebroids with anti-involution and trivial graded have been classified up to equivalence in [33] (see also [4, 5] for the possibly twisted case), whereas Morita-type results for deformation quantization algebras are obtained in [6, 8, 7] for real Poisson manifolds, and in [40] in the algebraic setting.

Acknowledgments. We express our gratitude to Masaki Kashiwara for useful discussions and for communicating Theorem 4.3.5 to us. We would like to thank the referee for pointing out a mistake in a previous version of this paper.

Convention. In this text, when dealing with categories and stacks, we will not mention any smallness condition (with respect to a given universe), leaving to the reader the task to make it precise when necessary.

1. NON-ABELIAN COHOMOLOGY

We are interested in classifying \mathcal{E} -algebroids and stacks of twisted \mathcal{E} -modules. Thanks to the existence of a canonical \mathcal{E} -algebroid, this amounts to classify stacks locally equivalent to a given one. To this end, we recall here some techniques of cohomology with values in a stack of 2-groups.

The classical reference for stacks is [15]. A quick introduction can be found in [11] and [22, chapter 19].

For stacks of 2-groups we refer to [3] (where the term *gr-stack* is used instead) and to [14, §1.4] for the strictly commutative case. See also [1] for an explicit description in terms of crossed modules. We follow the presentation of [33].

Let X be a topological space (or a site).

1.1. Stacks. A prestack \mathbf{C} on X is a lax presheaf of categories. It is lax in the sense that for a chain of three open subsets $W \subset V \subset U$ the restriction functor $\cdot|_W: \mathbf{C}(U) \rightarrow \mathbf{C}(W)$ coincides with the composition $\mathbf{C}(U) \xrightarrow{\cdot|_V} \mathbf{C}(V) \xrightarrow{\cdot|_W} \mathbf{C}(W)$ only up to an invertible transformation (such transformations need to satisfy a natural cocycle condition for chains of four open subsets).

For $\gamma, \gamma' \in \mathbf{C}(U)$, denote by $\mathcal{H}om_{\mathbf{C}}(\gamma, \gamma')$ the presheaf on U given by $U \supset V \mapsto \text{Hom}_{\mathbf{C}(V)}(\gamma|_V, \gamma'|_V)$. One says that \mathbf{C} is a separated prestack if $\mathcal{H}om_{\mathbf{C}}(\gamma, \gamma')$ is a sheaf for any γ, γ' . A stack on X is a separated prestack satisfying a natural descent condition, analogous to that for sheaves.

Given a stack \mathbf{C} , we denote by $\pi_0(\mathbf{C})$ the sheaf associated to the presheaf $X \supset U \mapsto \{\text{isomorphism classes of objects in } \mathbf{C}(U)\}$, and by

$\mathbf{Fct}(\mathbf{C}, \mathbf{C}')$, for another stack \mathbf{C}' , the stack whose objects on $U \subset X$ are functors from $\mathbf{C}|_U$ to $\mathbf{C}'|_U$ and whose morphisms are transformations.

Let $\varphi: Y \rightarrow X$ be a continuous map (or a morphism of sites). For \mathbf{D} a stack on Y and \mathbf{C} a stack on X , we denote by $\varphi_*\mathbf{D}$ and $\varphi^{-1}\mathbf{C}$ the stack-theoretical direct and inverse image, respectively. Recall that $\varphi^{-1}\mathbf{C}$ is the stack on Y associated to the separated prestack $\varphi^+\mathbf{C}$, defined on an open subset $V \subset Y$ by the category

$$\mathrm{Ob}(\varphi^+\mathbf{C}(V)) = \bigsqcup_{\substack{U \supset \varphi(V) \\ U \text{ open}}} \mathrm{Ob}(\mathbf{C}(U)),$$

$$\mathrm{Hom}_{\varphi^+\mathbf{C}(V)}(\gamma_U, \gamma_{U'}) = \Gamma(V; \varphi^{-1}\mathcal{H}om_{\mathbf{C}}(\gamma_U|_{U \cap U'}, \gamma_{U'}|_{U \cap U'})).$$

One checks that there is a natural equivalence (in fact, a 2-adjunction)

$$(1.1.1) \quad \varphi_*\mathbf{Fct}(\varphi^{-1}\mathbf{C}, \mathbf{D}) \approx \mathbf{Fct}(\mathbf{C}, \varphi_*\mathbf{D}).$$

Hence there are adjunction functors

$$\mathbf{C} \rightarrow \varphi_*\varphi^{-1}\mathbf{C}, \quad \varphi^{-1}\varphi_*\mathbf{D} \rightarrow \mathbf{D}.$$

By using the first adjunction, one gets an isomorphism of sheaves

$$(1.1.2) \quad \varphi^{-1}\pi_0(\mathbf{C}) \xrightarrow{\sim} \pi_0(\varphi^{-1}\mathbf{C}).$$

1.2. Stacks of 2-groups. Let \mathbf{C} be a stack on X . Denote by $\mathbf{Aut}(\mathbf{C})$ the substack of $\mathbf{Fct}(\mathbf{C}, \mathbf{C})$ whose objects are the auto-equivalences of \mathbf{C} , and whose morphisms are the *invertible* transformations. For $\mathcal{U} = \{U_i\}_{i \in I}$ an open cover of X , set

$$U_{ij} = U_i \cap U_j, \quad U_{ijk} = U_i \cap U_j \cap U_k, \quad \text{etc.}$$

Proposition A.1.1 for $\mathbf{C}_i = \mathbf{C}'_i = \mathbf{C}|_{U_i}$ describes how to patch objects and morphisms of $\mathbf{Aut}(\mathbf{C})$. With notations as in Proposition A.1.1, let $H^1(\mathcal{U}; \mathbf{Aut}(\mathbf{C}))$ be the pointed set of equivalence classes of pairs $(\mathbf{f}_{ij}, \mathbf{a}_{ijk})_{ijk \in I}$ satisfying the cocycle condition (A.1.1), modulo the coboundary relation given by pairs $(\mathbf{g}_i, \mathbf{b}_{ij})_{ij \in J}$ satisfying (A.1.2). As we recall in Remark A.1.2, an open cover \mathcal{V} finer than \mathcal{U} induces a well defined map $H^1(\mathcal{U}; \mathbf{Aut}(\mathbf{C})) \rightarrow H^1(\mathcal{V}; \mathbf{Aut}(\mathbf{C}))$. Hence one sets

$$(1.2.1) \quad H^1(X; \mathbf{Aut}(\mathbf{C})) = \varinjlim_{\mathcal{U}} H^1(\mathcal{U}; \mathbf{Aut}(\mathbf{C})).$$

By Proposition A.1.1 it follows immediately

Corollary 1.2.1. *The pointed set $H^1(X; \mathbf{Aut}(\mathbf{C}))$ is isomorphic to the pointed set of equivalence classes of stacks locally equivalent to \mathbf{C} .*

Let us recall how to make the construction (1.2.1) functorial.

A 2-group (also called a *gr-category* in [3, 1]) is a category endowed with a group structure both on morphisms and on objects. More precisely, a category \mathbf{G} is a 2-group if it is a groupoid (i.e. all morphisms are invertible) and it has a structure $(\mathbf{G}, \otimes, \mathbf{1})$ of monoidal category such

that each object $\gamma \in \mathbf{G}$ has a chosen (right) inverse. Functors of 2-groups and transformations between them are monoidal functors and monoidal transformations, respectively.

A stack \mathbf{G} is called a stack of 2-groups (a *gr-stack* in [3, 1]) if its categories of sections are 2-groups, its restrictions are functors of 2-groups and its transformations between restriction functors are monoidal. Functors of stacks of 2-groups are functors of monoidal stacks.

Recall that one sets $\pi_1(\mathbf{G}) = \mathcal{H}om_{\mathbf{G}}(\mathbf{1}, \mathbf{1})$. Both $\pi_0(\mathbf{G})$ and $\pi_1(\mathbf{G})$ are sheaves of groups, the latter being necessarily commutative. Any functor of stacks of 2-groups induces a group homomorphism at the level of π_1 and π_0 .

Example 1.2.2. For \mathcal{G} a sheaf of groups, denote by $\mathcal{G}[0]$ the stack obtained by trivially enriching \mathcal{G} with identity arrows, and by $\mathcal{G}[1]$ the stack of right \mathcal{G} -torsors. Then $\mathcal{G}[0]$ is a stack of 2-groups, and $\mathcal{G}[1]$ is a stack of 2-groups if and only if \mathcal{G} is commutative.

Another example of stack of 2-groups is given by $\mathbf{Aut}(\mathbf{C})$ for a stack \mathbf{C} . Let \mathbf{G} be a stack of 2-groups and \mathcal{U} an open cover of X . One can extend as follows the construction (1.2.1), where one should read “ \otimes ” instead of “ \circ ” in all diagrams in Appendix A.1.

A 1-cocycle with values in \mathbf{G} is a pair $(\gamma_{ij}, a_{ijk})_{ijk \in I}$ with $\gamma_{ij} \in \mathbf{G}(U_{ij})$ and $a_{ijk} \in \text{Hom}_{\mathbf{G}(U_{ijk})}(\gamma_{ik}, \gamma_{ij} \otimes \gamma_{jk})$ satisfying (A.1.1). Two such 1-cocycles $(\gamma_{ij}, a_{ijk})_{ijk \in I}$ and $(\gamma'_{ij}, a'_{ijk})_{ijk \in I}$ are cohomologous if there is a pair $(\delta_i, b_{ij})_{ij \in I}$ with $\delta_i \in \mathbf{G}(U_i)$ and $b_{ij} \in \text{Hom}_{\mathbf{G}(U_{ij})}(\gamma'_{ij} \otimes \delta_j, \delta_i \otimes \gamma_{ij})$ satisfying (A.1.2).

The first cohomology pointed set of \mathbf{G} on X is given by

$$H^1(X; \mathbf{G}) = \varinjlim_{\mathcal{U}} H^1(\mathcal{U}; \mathbf{G}),$$

where $H^1(\mathcal{U}; \mathbf{G})$ denotes the pointed set of equivalence classes of 1-cocycles on \mathcal{U} , modulo the relation of being cohomologous. One can also define the cohomology in degree 0 and -1 . This construction is functorial in the sense that short exact sequences of 2-groups induce long exact cohomology sequences (in a sense to be made precise). In particular, equivalent 2-groups have isomorphic cohomologies.

With notation as in Example 1.2.2 one has

$$(1.2.2) \quad H^1(X; \mathcal{G}[i]) \simeq H^{1+i}(X; \mathcal{G}) \quad \text{for } i = 0, 1,$$

where \mathcal{G} is assumed to be abelian if $i = 1$. Here, the pointed set $H^1(X; \mathcal{G})$ is defined by Čech cohomology and $H^2(X; \mathcal{G})$ is computed using hypercoverings.

1.3. Crossed modules. A crossed module is the data

$$\mathcal{G}^\bullet = (\mathcal{G}^{-1} \xrightarrow{d} \mathcal{G}^0, \delta)$$

of a complex of sheaves of groups and of a left action δ of \mathcal{G}^0 on \mathcal{G}^{-1} such that for any $f \in \mathcal{G}^0$ and $a \in \mathcal{G}^{-1}$

$$d \circ \delta(f) = \text{Ad}(f) \circ d, \quad \delta(d(a)) = \text{Ad}(a),$$

where $\text{Ad}(a)(b) = aba^{-1}$. A morphism of crossed modules is a morphism of complexes of sheaves of groups compatible with the left actions.

There is a functorial way of associating to a crossed module a stack of 2-groups as follows. For \mathcal{G}^\bullet a crossed module one denotes by $[\mathcal{G}^\bullet]$ the stack of 2-groups associated to the separated prestack whose objects on $U \subset X$ are sections $f \in \mathcal{G}^0(U)$ and whose morphisms $f \rightarrow f'$ are sections $a \in \mathcal{G}^{-1}(U)$ satisfying $f' = d(a)f$. Then $[\mathcal{G}^\bullet]$ is a stack of 2-groups, with monoidal structure given by $f \otimes g = fg$ at the level of objects and by $a \otimes b = a\delta(f)(b)$ at the level of morphisms, for $a: f \rightarrow f'$ and $b: g \rightarrow g'$.

One checks that there are isomorphisms of groups

$$\pi_i([\mathcal{G}^\bullet]) \simeq H^{-i}(\mathcal{G}^\bullet), \quad i = 0, 1$$

and, with notation and conventions as in Example 1.2.2, equivalences of stacks of 2-groups

$$[\mathcal{G}[i]] \approx \mathcal{G}[i], \quad i = 0, 1,$$

where the structure of crossed module on the complex $\mathcal{G}[i]$ (obtained by placing \mathcal{G} in $-i$ position) is the trivial one.

1.4. Strictly abelian crossed modules. Denote by $D^{[-1,0]}(\mathbb{Z}_X)$ the full subcategory of the derived category of sheaves of abelian groups whose objects have cohomology concentrated in degree -1 and 0 . Consider a complex of abelian groups $\mathcal{F}^\bullet \in C^{[-1,0]}(\mathbb{Z}_X)$ as a crossed module with trivial left action. Then the functor $\mathcal{F}^\bullet \mapsto [\mathcal{F}^\bullet]$ factorizes through $D^{[-1,0]}(\mathbb{Z}_X)$ and one has

$$(1.4.1) \quad H^1(X; \mathcal{F}^\bullet) = H^1(X; [\mathcal{F}^\bullet]).$$

Let $\psi: X \rightarrow Y$ be a continuous map (or a morphism of sites). The inverse and direct image of stacks of 2-groups are again stacks of 2-groups, and one has

$$(1.4.2) \quad \psi^{-1}[\mathcal{G}^\bullet] \approx [\psi^{-1}\mathcal{G}^\bullet] \quad \psi_*[\mathcal{F}^\bullet] \approx [\tau_{\leq 0} R\psi_* \mathcal{F}^\bullet],$$

where $\tau_{\leq 0}$ is the truncation functor. In particular, for a sheaf of abelian groups \mathcal{F} , one gets

$$(1.4.3) \quad \pi_i(\psi_*(\mathcal{F}[1])) \simeq R^{1-i}\psi_* \mathcal{F}, \quad i = 0, 1.$$

2. ALGEBROIDS

Mitchell [30] showed how algebras can be replaced by linear categories. Similarly, sheaves of algebras can be replaced by linear stacks. An algebroid is a linear stack locally equivalent to an algebra. This notion, already implicit in [17], was introduced in [26] and developed in [12] (see

also [23, §2.1] and [10]). It is the linear analogue of the notion of gerbe from [15]: an algebroid is to a gerbe as an algebra is to a group.

Let X be a topological space (or a site), and \mathcal{R} a sheaf of commutative rings on X .

2.1. Linear stacks. A stack \mathcal{C} on X is called \mathcal{R} -linear (\mathcal{R} -stack, for short) if for any $\gamma, \gamma' \in \mathcal{C}(U)$ the sheaf $\mathcal{H}om_{\mathcal{C}}(\gamma, \gamma')$ is endowed with an $\mathcal{R}|_U$ -module structure compatible with composition. In particular, $\mathcal{E}nd_{\mathcal{C}}(\gamma)$ has an $\mathcal{R}|_U$ -algebra structure with product given by composition. A functor between \mathcal{R} -linear stacks is called \mathcal{R} -linear (\mathcal{R} -functor, for short) if it is \mathcal{R} -linear at the level of morphisms, while no linearity conditions are required on transformations.

One says that two \mathcal{R} -stacks are equivalent if they are equivalent through an \mathcal{R} -functor. This implies that the quasi-inverse is also an \mathcal{R} -functor. We denote by $\approx_{\mathcal{R}}$ this equivalence relation.

The center $Z(\mathcal{C})$ of an \mathcal{R} -stack \mathcal{C} is the sheaf of endo-transformations of the identity functor $\text{id}_{\mathcal{C}}$. It has a natural structure of sheaf of commutative \mathcal{R} -algebras. Note that a stack \mathcal{C} is \mathcal{R} -linear if and only if it is \mathbb{Z} -linear and its center is an \mathcal{R} -algebra.

If \mathcal{C} is an \mathcal{R} -stack, then its opposite stack \mathcal{C}^{op} is again \mathcal{R} -linear. For \mathcal{D} another \mathcal{R} -stack, $\text{Fct}_{\mathcal{R}}(\mathcal{C}, \mathcal{D})$ denotes the full substack of $\text{Fct}(\mathcal{C}, \mathcal{D})$ whose objects are \mathcal{R} -functors, and is itself an \mathcal{R} -stack. The tensor product $\mathcal{C} \otimes_{\mathcal{R}} \mathcal{D}$ is the \mathcal{R} -stack associated with the prestack $U \mapsto \mathcal{C}(U) \otimes_{\mathcal{R}(U)} \mathcal{D}(U)$ whose objects are pairs in $\mathcal{C}(U) \times \mathcal{D}(U)$, with morphisms

$$\text{Hom}_{\mathcal{C}(U) \otimes_{\mathcal{R}(U)} \mathcal{D}(U)}((\gamma, \delta), (\gamma', \delta')) = \text{Hom}_{\mathcal{C}(U)}(\gamma, \gamma') \otimes_{\mathcal{R}(U)} \text{Hom}_{\mathcal{D}(U)}(\delta, \delta').$$

Lemma 2.1.1. *If \mathcal{R} is an \mathcal{S} -algebra and \mathcal{E} an \mathcal{S} -stack, then*

$$\text{Fct}_{\mathcal{S}}(\mathcal{C} \otimes_{\mathcal{R}} \mathcal{D}, \mathcal{E}) \approx_{\mathcal{R}} \text{Fct}_{\mathcal{R}}(\mathcal{C}, \text{Fct}_{\mathcal{S}}(\mathcal{D}, \mathcal{E})).$$

(This is in fact a 2-adjunction.)

Let $\varphi: Y \rightarrow X$ be a continuous map (or a morphism of sites). Then $\varphi^{-1}\mathcal{C}$ is $\varphi^{-1}\mathcal{R}$ -linear and there is a $\varphi^{-1}\mathcal{R}$ -equivalence

$$\varphi^{-1}(\mathcal{C} \otimes_{\mathcal{R}} \mathcal{D}) \approx \varphi^{-1}\mathcal{C} \otimes_{\varphi^{-1}\mathcal{R}} \varphi^{-1}\mathcal{D}.$$

If \mathcal{E} is a $\varphi^{-1}\mathcal{R}$ -stack, then $\varphi_*\mathcal{E}$ is \mathcal{R} -linear and there is an \mathcal{R} -functor

$$(2.1.1) \quad \varphi_*\mathcal{E} \otimes_{\mathcal{R}} \varphi_*\mathcal{F} \rightarrow \varphi_*(\mathcal{E} \otimes_{\varphi^{-1}\mathcal{R}} \mathcal{F}).$$

2.2. Modules over a linear stack. Denote by $\text{Mod}(\mathcal{R})$ the category of \mathcal{R} -modules and by $\text{Mod}(\mathcal{R})$ the corresponding \mathcal{R} -stack given by $U \mapsto \text{Mod}(\mathcal{R}|_U)$.

For \mathcal{C} an \mathcal{R} -stack, the \mathcal{R} -stack of (left) \mathcal{C} -modules is defined by

$$(2.2.1) \quad \text{Mod}(\mathcal{C}) = \text{Fct}_{\mathcal{R}}(\mathcal{C}, \text{Mod}(\mathcal{R})).$$

(It follows from Lemma 2.3.4 that this definition does not depend on the base ring.)

The contravariant 2-functor $\mathbf{Mod}(\cdot)$ is defined as follows. On objects, it is given by (2.2.1). Consider the diagram

$$\begin{array}{ccc} & \xrightarrow{f} & \\ \text{C} & \Downarrow \text{d} & \text{D} \xrightarrow{\mathcal{N}} \mathbf{Mod}(\mathcal{R}), \\ & \xrightarrow{f'} & \end{array}$$

where f, f' and \mathcal{N} are \mathcal{R} -functors, and d a transformation. To the \mathcal{R} -functor $f: \text{C} \rightarrow \text{D}$ one associates the \mathcal{R} -functor

$$\mathbf{Mod}(f): \mathbf{Mod}(\text{D}) \rightarrow \mathbf{Mod}(\text{C}), \quad \mathcal{N} \mapsto \mathcal{N} \circ f,$$

and to the transformation $d: f \rightarrow f'$ one associates the transformation,

$$\mathbf{Mod}(d): \mathbf{Mod}(f) \rightarrow \mathbf{Mod}(f'),$$

such that $\mathbf{Mod}(d)(\mathcal{N}) = \text{id}_{\mathcal{N}} \bullet d$ is the morphism associated to $\mathcal{N} \in \mathbf{Mod}(\text{D})$, where \bullet denotes the horizontal composition of transformations. In other words, for $\gamma \in \text{C}$ one has $\mathbf{Mod}(d)(\mathcal{N})(\gamma) = \mathcal{N}(d(\gamma))$ as morphisms from $\mathcal{N}(f(\gamma))$ to $\mathcal{N}(f'(\gamma))$ in $\mathbf{Mod}(\mathcal{R})$. We use the notation

$$(2.2.2) \quad \mathfrak{f}(\cdot) = \mathbf{Mod}(f).$$

2.3. Algebras as stacks. Let \mathcal{A} be a sheaf of \mathcal{R} -algebras. Denote by \mathcal{A}^{op} the opposite algebra, by $\mathbf{Mod}(\mathcal{A})$ the \mathcal{R} -stack of left \mathcal{A} -modules and by $\mathcal{H}om_{\mathcal{A}}(\cdot, \cdot) = \mathcal{H}om_{\mathbf{Mod}(\mathcal{A})}(\cdot, \cdot)$ the internal hom-functor.

Denote by \mathcal{A}^+ the full substack of $\mathbf{Mod}(\mathcal{A}^{\text{op}})$ whose objects are locally free right \mathcal{A} -modules of rank one. For any $\mathcal{N} \in \mathcal{A}^+(U)$ there is an $\mathcal{R}|_U$ -algebra isomorphism $\mathcal{E}nd_{\mathcal{A}^+}(\mathcal{N}) \simeq \mathcal{A}|_U$. Note that the stack \mathcal{A}^+ has a canonical global object given by \mathcal{A} itself with its structure of right \mathcal{A} -module. In particular, the sheaf $\pi_0(\mathcal{A}^+)$ is the singleton-valued constant sheaf.

For $f: \mathcal{A} \rightarrow \mathcal{B}$ an \mathcal{R} -algebra morphism, denote by $f^+: \mathcal{A}^+ \rightarrow \mathcal{B}^+$ the \mathcal{R} -functor induced by the extension of scalars $(\cdot) \otimes_{\mathcal{A}} \mathcal{B}$. We thus have a functor between the stack of \mathcal{R} -algebras and that of \mathcal{R} -stacks

$$(\cdot)^+: \mathcal{R}\text{-Alg}_X \rightarrow \mathcal{R}\text{-Stk}_X.$$

Remark 2.3.1. Let $\mathcal{A}^{“+”}$ be the separated prestack $U \mapsto \mathcal{A}(U)^+$, where $\mathcal{A}(U)^+$ denotes the $\mathcal{R}(U)$ -category with one object and sections of $\mathcal{A}(U)$ as its endomorphisms. By Yoneda lemma (see §3.1), the stack associated to $\mathcal{A}^{“+”}$ is \mathcal{R} -equivalent to \mathcal{A}^+ .

The stack $\mathcal{R}\text{-Stk}_X$ is naturally upgraded to a 2-stack by considering transformations of functors. Note that for any two \mathcal{R} -algebras \mathcal{A} and \mathcal{B} on $U \subset X$ and any \mathcal{R} -functor $f: \mathcal{A}^+ \rightarrow \mathcal{B}^+$ there exist a cover $\mathcal{U} = \{U_i\}_{i \in I}$ of U and morphisms of \mathcal{R} -algebras $f_i: \mathcal{A}|_{U_i} \rightarrow \mathcal{B}|_{U_i}$ such that $f|_{U_i} \simeq f_i^+$.

Definition 2.3.2. One says that an \mathcal{R} -stack \mathbf{C} is equivalent to an \mathcal{R} -algebra \mathcal{A} if $\mathbf{C} \approx_{\mathcal{R}} \mathcal{A}^+$.

In Proposition 2.6.2 we characterize the condition of equivalence between algebras.

Recall that a stack \mathbf{C} is non-empty if it has at least one global object, and it is locally connected by isomorphisms if any two objects $\gamma, \gamma' \in \mathbf{C}(U)$ are locally isomorphic. If \mathbf{C} is \mathcal{R} -linear, this amounts to ask that the sheaf $\mathcal{H}om_{\mathbf{C}}(\gamma, \gamma')$ is a locally free $\mathcal{E}nd_{\mathbf{C}}(\gamma')$ -module of rank one.

Lemma 2.3.3. *An \mathcal{R} -stack \mathbf{C} is equivalent to an \mathcal{R} -algebra if and only if it is non-empty and locally connected by isomorphisms.*

Proof. One implication is clear. Suppose that \mathbf{C} is non-empty and let $\gamma \in \mathbf{C}(X)$. Then the fully faithful functor $\mathcal{E}nd_{\mathbf{C}}(\gamma)^+ \rightarrow \mathbf{C}$ is an equivalence if and only if \mathbf{C} is locally connected by isomorphisms. \square

Let \mathbf{C} be an \mathcal{R} -stack. For $\mathcal{N} \in \mathcal{R}^+$ and $\gamma \in \mathbf{C}$, one defines $\mathcal{N} \otimes_{\mathcal{R}} \gamma \in \mathbf{C}$ as the representative of $\mathcal{N} \otimes_{\mathcal{R}} \mathcal{H}om_{\mathbf{C}}(\cdot, \gamma) \in \mathbf{Mod}(\mathbf{C}^{\text{op}})$. Then one has \mathcal{R} -equivalences

$$\begin{aligned} \mathcal{R}^+ \otimes_{\mathcal{R}} \mathbf{C} &\approx_{\mathcal{R}} \mathbf{C}, & (\mathcal{N}, \gamma) &\mapsto \mathcal{N} \otimes_{\mathcal{R}} \gamma, \\ \mathbf{C} &\approx_{\mathcal{R}} \mathbf{Fct}_{\mathcal{R}}(\mathcal{R}^+, \mathbf{C}), & \gamma &\mapsto (\cdot) \otimes_{\mathcal{R}} \gamma. \end{aligned}$$

Lemma 2.3.4. *The definition (2.2.1) of stack of \mathbf{C} -modules does not depend on the base ring \mathcal{R} .*

Proof. It follows from Lemma 2.1.1 for $\mathcal{S} = \mathbb{Z}_X$, $\mathbf{D} = \mathcal{R}^+$ and $\mathbf{E} = \mathbf{Mod}(\mathbb{Z}_X)$ that

$$\mathbf{Fct}_{\mathcal{R}}(\mathbf{C}, \mathbf{Mod}(\mathcal{R})) \approx_{\mathcal{R}} \mathbf{Fct}_{\mathbb{Z}_X}(\mathbf{C}, \mathbf{Mod}(\mathbb{Z}_X)),$$

where we use the equivalence $\mathbf{Fct}_{\mathbb{Z}_X}(\mathcal{R}^+, \mathbf{Mod}(\mathbb{Z}_X)) \approx_{\mathcal{R}} \mathbf{Mod}(\mathcal{R})$. \square

2.4. Compatibility. Let \mathcal{A} and \mathcal{B} be two \mathcal{R} -algebras, and $\varphi: Y \rightarrow X$ a continuous map (or a morphism of sites). There are an \mathcal{R} -algebra isomorphism

$$Z(\mathcal{A}) \xrightarrow{\sim} Z(\mathcal{A}^+), \quad a \mapsto (\mathcal{N} \rightarrow \mathcal{N}; n \mapsto an),$$

and \mathcal{R} -equivalences

$$\begin{aligned} (\mathcal{A}^+)^{\text{op}} &\approx_{\mathcal{R}} (\mathcal{A}^{\text{op}})^+, & \mathcal{N} &\mapsto \mathcal{H}om_{\mathcal{A}^{\text{op}}}(\mathcal{N}, \mathcal{A}), \\ \mathbf{Mod}(\mathcal{A}) &\approx_{\mathcal{R}} \mathbf{Mod}(\mathcal{A}^+), & \mathcal{M} &\mapsto (\cdot) \otimes_{\mathcal{A}} \mathcal{M}, \\ \mathcal{A}^+ \otimes_{\mathcal{R}} \mathcal{B}^+ &\approx_{\mathcal{R}} (\mathcal{A} \otimes_{\mathcal{R}} \mathcal{B})^+, & (\mathcal{N}, \mathcal{Q}) &\mapsto \mathcal{N} \otimes_{\mathcal{R}} \mathcal{Q} \\ \varphi^{-1} \mathcal{A}^+ &\approx_{\mathcal{R}} (\varphi^{-1} \mathcal{A})^+, & \mathcal{N} &\mapsto \varphi^{-1} \mathcal{N}. \end{aligned}$$

For $f, f': \mathcal{A} \rightarrow \mathcal{B}$ two \mathcal{R} -algebra morphisms, the sections on $U \subset X$ of the sheaf $\mathcal{H}om_{\mathbf{Fct}_{\mathcal{R}}(\mathcal{A}^+, \mathcal{B}^+)}(f^+, f'^+)$ are given by

$$(2.4.1) \quad \{b \in \mathcal{B}(U): bf(a) = f'(a)b \text{ for each } a \in \mathcal{A}(V) \text{ and } V \subset U\},$$

with composition of transformations given by the product in \mathcal{B} .

For \mathcal{N} a left \mathcal{B} -module, with notation (2.2.2) we set

$$(2.4.2) \quad {}_f\mathcal{N} = {}_{f^+}\mathcal{N}$$

the associated left \mathcal{A} -module. By (2.4.1), the morphism of \mathcal{A} -modules associated to a transformation $b: f^+ \rightarrow f'^+$ is given by

$${}_f\mathcal{N} \rightarrow {}_{f'}\mathcal{N}; n \mapsto bn.$$

2.5. Algebroids. Recall from Lemma 2.3.3 that an \mathcal{R} -stack is equivalent to an \mathcal{R} -algebra if and only if it is non-empty and is locally connected by isomorphisms.

Definition 2.5.1. An \mathcal{R} -algebroid is an \mathcal{R} -stack which is locally non-empty and locally connected by isomorphisms.

In other words, an \mathcal{R} -algebroid is an \mathcal{R} -stack \mathbf{A} which is locally equivalent to an algebra. It is globally an algebra if and only if it has a global object¹. Note also that an \mathcal{R} -stack is an \mathcal{R} -algebroid if and only if its substack with the same objects and only invertible morphisms is a gerbe.

The stack $\mathbf{Mod}(\mathbf{A})$ is an example of stack of twisted sheaves, i.e. it is a stack locally equivalent to a stack of modules over an algebra (see [22, 11]). A cocyclic description of algebroids and of their modules is recalled in Appendix A.2 and A.3.

Note that the existence of an \mathcal{R} -functor $\mathcal{R}^+ \rightarrow \mathbf{A}$ is equivalent to the existence of a global object for \mathbf{A} . In this case there is a forgetful functor

$$\mathbf{Mod}(\mathbf{A}) \rightarrow \mathbf{Mod}(\mathcal{R}).$$

Lemma 2.5.2. *An \mathcal{R} -stack \mathbf{C} on X is an algebroid if and only if $\pi_0(\mathbf{C}) \simeq \{\text{pt}\}_X$, the singleton-valued constant sheaf.*

It follows from (1.1.2) that inverse images of algebroids are algebroids.

Let \mathbf{C} be an \mathcal{R} -stack. Then for any \mathcal{R} -algebroid \mathbf{A} one has

$$\pi_0(\mathbf{A} \otimes_{\mathcal{R}} \mathbf{C}) \simeq \pi_0(\mathbf{C}).$$

In particular, the tensor product of algebroids is an algebroid.

Definition 2.5.3. (i) Let \mathcal{A} be an \mathcal{R} -algebra. An \mathcal{R} -twisted form of \mathcal{A} is an \mathcal{R} -algebroid locally \mathcal{R} -equivalent to \mathcal{A} .
(ii) An invertible \mathcal{R} -algebroid is an \mathcal{R} -twisted form of \mathcal{R} .

Note that any \mathcal{R} -functor between invertible \mathcal{R} -algebroids is an equivalence, since it is locally isomorphic to the identity functor of \mathcal{R}^+ .

¹If the category $\mathbf{A}(U)$ has a zero object for $U \subset X$, then $\mathbf{A}|_U \approx 0^+$, where 0 denotes the ring with $1 = 0$. In particular, except for the case 0^+ , algebroids are not stacks of additive categories.

If \mathcal{C} is an invertible \mathcal{R} -algebroid, then $\mathcal{R} \xrightarrow{\sim} Z(\mathcal{C})$ and for any \mathcal{R} -stack \mathcal{D} there is an \mathcal{R} -equivalence

$$\mathcal{C}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{D} \approx_{\mathcal{R}} \text{Fct}_{\mathcal{R}}(\mathcal{C}, \mathcal{D}), \quad (\gamma, \delta) \mapsto \text{Hom}_{\mathcal{C}}(\gamma, \cdot) \otimes_{\mathcal{R}} \delta.$$

In particular, the set of \mathcal{R} -equivalence classes of invertible \mathcal{R} -algebroids is a group, with multiplication given by $\otimes_{\mathcal{R}}$ and inverse given by $(\cdot)^{\text{op}}$.

For an \mathcal{R} -stack \mathcal{C} , denote by $\text{Aut}_{\mathcal{R}}(\mathcal{C})$ the full substack of $\text{Aut}(\mathcal{C})$ whose objects are \mathcal{R} -equivalences. By Corollary 1.2.1, the cohomology $H^1(X; \text{Aut}_{\mathcal{R}}(\mathcal{A}^+))$ classifies \mathcal{R} -equivalence classes of \mathcal{R} -twisted forms of \mathcal{A} . In terms of crossed modules (cf. Section 1.3), one has

$$(2.5.1) \quad \text{Aut}_{\mathcal{R}}(\mathcal{A}^+) \approx [(\mathcal{A}^{\times} \xrightarrow{\text{Ad}} \text{Aut}_{\mathcal{R}\text{-Alg}_X}(\mathcal{A}), \delta)],$$

where $\delta(f)(a) = f(a)$ and $\text{Ad}(a)(b) = aba^{-1}$. In particular, $\text{Aut}_{\mathcal{R}}(\mathcal{R}^+) \approx \mathcal{R}^{\times}[1]$ and (1.2.2) implies

Lemma 2.5.4. *The group of \mathcal{R} -equivalence classes of invertible \mathcal{R} -algebroids is isomorphic to $H^2(X; \mathcal{R}^{\times})$.*

2.6. Inner forms. Let \mathcal{A} be a central \mathcal{R} -algebra, i.e. $Z(\mathcal{A}) = \mathcal{R}$. (If \mathcal{A} is not central, the following discussion still holds by replacing \mathcal{R} with $Z(\mathcal{A})$.)

Denote by $\mathcal{I}nn(\mathcal{A})$ the sheaf theoretical image of the group morphism $\text{Ad}: \mathcal{A}^{\times} \rightarrow \text{Aut}_{\mathcal{R}\text{-Alg}_X}(\mathcal{A})$. Its sections are the inner automorphisms of \mathcal{A} , i.e. automorphisms locally of the form $\text{Ad}(a)$ for some $a \in \mathcal{A}^{\times}$. Recall that an \mathcal{R} -algebra \mathcal{B} is called an inner form of \mathcal{A} if there exists an open cover $\{U_i\}_{i \in I}$ of X and ring isomorphisms $f_i: \mathcal{A}|_{U_i} \rightarrow \mathcal{B}|_{U_i}$ such that $f_j^{-1}f_i \in \mathcal{I}nn(\mathcal{A}|_{U_{ij}})$.

Examples of inner forms are given by Azumaya algebras and rings of twisted differential operators (see for example [11] for more details).

Let \mathcal{B} be an \mathcal{R} -algebra. Denote by $\mathbf{E}_{\mathcal{A}, \mathcal{B}} \subset \text{Fct}_{\mathcal{R}}(\mathcal{A}^+, \mathcal{B}^+)$ the full substack of \mathcal{R} -equivalences. Note that $\mathbf{E}_{\mathcal{A}, \mathcal{B}}^{\text{op}} \approx_{\mathcal{R}} \mathbf{E}_{\mathcal{B}, \mathcal{A}}$.

Lemma 2.6.1. *\mathcal{B} is an inner form of \mathcal{A} if and only if $\mathbf{E}_{\mathcal{A}, \mathcal{B}}$ is an \mathcal{R} -algebroid.*

Proof. Since \mathcal{R} -equivalences $\mathcal{A}^+ \xrightarrow{\sim} \mathcal{B}^+$ are locally given by \mathcal{R} -algebra isomorphisms $\mathcal{A} \xrightarrow{\sim} \mathcal{B}$, it follows that $\mathbf{E}_{\mathcal{A}, \mathcal{B}}$ is locally non-empty if and only if \mathcal{B} is locally isomorphic to \mathcal{A} .

Let $f, f': \mathcal{A} \rightarrow \mathcal{B}$ be \mathcal{R} -algebra isomorphisms. Setting $a = f^{-1}(b)$ in (2.4.1), the invertible transformations from f^+ to f'^+ are given by

$$\{a \in \mathcal{A}^{\times} : f^{-1}f' = \text{Ad}(a)\},$$

hence $\mathbf{E}_{\mathcal{A}, \mathcal{B}}$ is an \mathcal{R} -algebroid if and only if \mathcal{B} is an inner form of \mathcal{A} . \square

Since $\mathcal{E}nd_{\mathbf{E}_{\mathcal{A}, \mathcal{B}}}(f^+) = \mathcal{R}$, if \mathcal{B} is an inner form of \mathcal{A} it follows that $\mathbf{E}_{\mathcal{A}, \mathcal{B}}$ is an invertible \mathcal{R} -algebroid and $\mathbf{E}_{\mathcal{A}, \mathcal{B}} \otimes_{\mathcal{R}} \mathcal{A}^+ \approx_{\mathcal{R}} \mathcal{B}^+$. In particular, one gets an equivalence of stacks of 2-groups $\text{Aut}_{\mathcal{R}}(\mathcal{A}^+) \approx \text{Aut}_{\mathcal{R}}(\mathcal{B}^+)$.

Consider the non-abelian exact sequence

$$H^1(X; \mathcal{A}^\times) \xrightarrow{b} H^1(X; \mathcal{I}nn(\mathcal{A})) \xrightarrow{c} H^2(X; \mathcal{R}^\times)$$

induced by the short exact sequence

$$1 \rightarrow \mathcal{R}^\times \rightarrow \mathcal{A}^\times \xrightarrow{\text{Ad}} \mathcal{I}nn(\mathcal{A}) \rightarrow 1.$$

For \mathcal{B} an inner form of \mathcal{A} and \mathcal{P} a locally free \mathcal{A}^{op} -module of rank one, denote by $[\mathcal{B}]$ and $[\mathcal{P}]$ the associated cohomology classes in $H^1(X; \mathcal{I}nn(\mathcal{A}))$ and $H^1(X; \mathcal{A}^\times)$ respectively. Then $b[\mathcal{P}] = [\mathcal{E}nd_{\mathcal{A}^{\text{op}}}(\mathcal{P})]$ and $c([\mathcal{B}]) = [\mathbf{E}_{\mathcal{A}, \mathcal{B}}]$.

Proposition 2.6.2. *The following conditions are equivalent.*

- (i) *The stacks \mathcal{A}^+ and \mathcal{B}^+ are \mathcal{R} -equivalent.*
- (ii) *There exists a locally free \mathcal{A}^{op} -module \mathcal{P} of rank one such that $\mathcal{B} \simeq \mathcal{E}nd_{\mathcal{A}^{\text{op}}}(\mathcal{P})$.*
- (iii) *\mathcal{B} is an inner form of \mathcal{A} and $c([\mathcal{B}]) = 1$.*

Proof. (i) \Rightarrow (ii)² Let $\mathbf{g}: \mathcal{B}^+ \rightarrow \mathcal{A}^+$ be an \mathcal{R} -equivalence. Recall that $\mathcal{B}^+ \subset \text{Mod}(\mathcal{B}^{\text{op}})$ is the substack of locally free modules of rank one. Let β be the canonical global object of \mathcal{B}^+ , and set $\mathcal{P} = \mathbf{g}(\beta)$. Then \mathcal{B} is isomorphic to $\mathcal{E}nd_{\mathcal{A}^{\text{op}}}(\mathcal{P})$.

(ii) \Rightarrow (iii) \mathcal{B} is clearly an inner form of \mathcal{A} and \mathcal{P} has a structure of $\mathcal{A}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{B}$ -module by the isomorphism $\mathcal{B} \xrightarrow{\sim} \mathcal{E}nd_{\mathcal{A}^{\text{op}}}(\mathcal{P})$. Then $(\cdot) \otimes_{\mathcal{B}} \mathcal{P}$ gives a global object of $\mathbf{E}_{\mathcal{B}, \mathcal{A}}$ and $c([\mathcal{B}]) = [\mathbf{E}_{\mathcal{B}, \mathcal{A}}^{\text{op}}] = 1$.

(i) \Leftarrow (iii) By Lemma 2.6.1 follows that $c([\mathcal{B}]) = 1$ if and only if $\mathbf{E}_{\mathcal{A}, \mathcal{B}}$ has a global object.

□

3. MORITA THEORY FOR LINEAR STACKS

Morita theory classically deals with modules over algebras. It is extended to modules over linear categories in [29, 31] and to stacks of modules over sheaves of algebras in [22, Chapter 19] (see also [11]). Here, we summarize these extensions by considering stacks of modules over linear stacks, and in particular over algebroids.

Let X be a topological space (or a site), and \mathcal{R} a sheaf of commutative rings on X .

²The equivalence between (i) and (ii) can also be deduced from Corollary 3.3.8.

3.1. Yoneda embedding. Recall that a category is called (co)complete if it admits small (co)limits. A prestack \mathbf{C} on X is called (co)complete if the categories $\mathbf{C}(U)$ are (co)complete for each $U \subset X$, and the restriction functors commute with (co)limits.

A prestack \mathbf{C} on X is called a proper stack (see [21, 36]) if it is separated, cocomplete, and if for each inclusion of open subsets $v: V \hookrightarrow U$, the restriction functors $\mathbf{C}(v) = (\cdot)|_V$ admits a fully faithful left adjoint

$$v_!: \mathbf{C}(V) \rightarrow \mathbf{C}(U),$$

called zero-extension, such that for a diagram of open inclusions

$$\begin{array}{ccc} V \cap W & \xrightarrow{v'} & W \\ w' \downarrow & & \downarrow w \\ V & \xrightarrow{v} & U, \end{array}$$

the natural transformation $v'_! \circ \mathbf{C}(w') \rightarrow \mathbf{C}(w) \circ v_!$ is an isomorphism.

Lemma 3.1.1. *Let \mathbf{C} be a proper stack. For $\gamma \in \mathbf{C}(V)$ and $\gamma' \in \mathbf{C}(U)$ there is an isomorphism of $\mathcal{R}|_U$ -modules*

$$v_* \mathcal{H}om_{\mathbf{C}|_V}(\gamma, \gamma'|_V) \simeq \mathcal{H}om_{\mathbf{C}|_U}(v_! \gamma, \gamma').$$

Recall that proper stacks are stacks.

Lemma 3.1.2. *For any \mathcal{R} -stack \mathbf{C} , the \mathcal{R} -stack $\mathbf{Mod}(\mathbf{C})$ is proper and complete.*

Proof. Recall first that $\mathbf{Mod}(\mathcal{R})$ is complete and cocomplete. It is also proper. In fact, for $v: V \hookrightarrow U$ an open inclusion, the restriction functor of $\mathbf{Mod}(\mathcal{R})$ coincides with the sheaf-theoretical pull-back v^{-1} . This admits the direct image functor v_* as right adjoint, and the zero-extension functor $v_!$ as a left adjoint.

The statement follows, as $\mathbf{Mod}(\mathbf{C}) = \mathbf{Fct}_{\mathcal{R}}(\mathbf{C}, \mathbf{Mod}(\mathcal{R}))$ inherits the properties and structures of $\mathbf{Mod}(\mathcal{R})$. For example, for $v: V \hookrightarrow U$ an inclusion of open subsets, the functor $v_!: \mathbf{Mod}(\mathbf{C}|_V) \rightarrow \mathbf{Mod}(\mathbf{C}|_U)$ is given by $(v_! \mathcal{M})(\gamma) = u_!(\mathcal{M}(\gamma|_{V \cap W}))$, where $\mathcal{M}: \mathbf{C}|_V \rightarrow \mathbf{Mod}(\mathcal{R}|_V)$ is a $\mathbf{C}|_V$ -module, $W \subset U$ is an open subset, $\gamma \in \mathbf{C}(W)$, and $u: V \cap W \rightarrow U$ is the embedding. \square

Let \mathbf{C} be an \mathcal{R} -stack. The (linear) Yoneda embedding is the full and faithful \mathcal{R} -functor

$$(3.1.1) \quad \mathbf{Y}_{\mathbf{C}}: \mathbf{C}^{\text{op}} \rightarrow \mathbf{Mod}(\mathbf{C}), \quad \gamma \mapsto \mathcal{H}om_{\mathbf{C}}(\gamma, \cdot)$$

whose essential image are the functors $\mathbf{C} \rightarrow \mathbf{Mod}(\mathcal{R})$ which are representable. In analogy with the case $\mathbf{C} = \mathcal{A}^+$ for \mathcal{A} an \mathcal{R} -algebra, a module $\mathcal{M} \in \mathbf{Mod}(\mathbf{C})$ which is representable is called locally free of rank one.

As in the classical case, the full faithfulness of (3.1.1) follows from

Lemma 3.1.3. *For $\mathcal{M} \in \text{Mod}(\mathcal{C})$ there is an isomorphism of \mathcal{C} -modules*

$$\mathcal{M}(\cdot) \simeq \text{Hom}_{\text{Mod}(\mathcal{C})}(\mathcal{Y}_{\mathcal{C}}(\cdot), \mathcal{M}).$$

Let \mathcal{C} be an \mathcal{R} -stack. We use the notation

$$(3.1.2) \quad \mathcal{C} \in \text{Mod}(\mathcal{C}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{C})$$

for the canonical object $\text{Hom}_{\mathcal{C}}(\cdot, \cdot)$. This corresponds to the Yoneda embedding $\mathcal{Y}_{\mathcal{C}}$ via the equivalence induced by Lemma 2.1.1

$$\text{Mod}(\mathcal{C}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{C}) \approx_{\mathcal{R}} \text{Fct}_{\mathcal{R}}(\mathcal{C}^{\text{op}}, \text{Mod}(\mathcal{C})).$$

If $\mathcal{C} = \mathcal{A}^+$, the object in (3.1.2) coincides with \mathcal{A} , considered as a bimodule over itself. If \mathcal{C} is an invertible \mathcal{R} -algebroid, then $\mathcal{C}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{C} \approx_{\mathcal{R}} \mathcal{R}^+$ and \mathcal{C} is isomorphic to \mathcal{R} as a bimodule over itself.

Lemma 3.1.4. *For \mathcal{C} an \mathcal{R} -stack, there is a natural isomorphism of \mathcal{R} -algebras $Z(\mathcal{C}) \simeq Z(\text{Mod}(\mathcal{C}))$.*

Proof. Recall that the 2-functor $\text{Mod}(\cdot)$ defines a morphism of \mathcal{R} -algebras $Z(\mathcal{C}) \rightarrow Z(\text{Mod}(\mathcal{C}))$, sending a transformation \mathbf{d} of $\text{id}_{\mathcal{C}}$ to the transformation of $\text{id}_{\text{Mod}(\mathcal{C})}$ whose morphism associated to $\mathcal{N} \in \text{Mod}(\mathcal{C})$ is $\text{id}_{\mathcal{N}} \bullet \mathbf{d}$. (Here \bullet denotes the horizontal composition of transformations). Hence we get a commutative diagram

$$\begin{array}{ccc} Z(\mathcal{C}) & \xrightarrow{\quad} & Z(\text{Mod}(\mathcal{C})) \\ \parallel & & \downarrow \cdot \text{id}_{\mathcal{Y}_{\mathcal{C}}} \\ Z(\mathcal{C}^{\text{op}}) & \xrightarrow{\text{id}_{\mathcal{Y}_{\mathcal{C}}} \bullet \cdot} & \mathcal{E}nd_{\text{Fct}_{\mathcal{R}}(\mathcal{C}^{\text{op}}, \text{Mod}(\mathcal{C}))}(\mathcal{Y}_{\mathcal{C}}) \end{array}$$

where $\text{id}_{\mathcal{Y}_{\mathcal{C}}} \bullet \cdot$ is an isomorphism, since (3.1.1) is fully faithful, and $\cdot \bullet \text{id}_{\mathcal{Y}_{\mathcal{C}}}$ is injective, since by Lemma 3.1.3 a transformation \mathbf{c} of $\text{id}_{\text{Mod}(\mathcal{C})}$ is completely determined by $\mathbf{c} \bullet \text{id}_{\mathcal{Y}_{\mathcal{C}}}$. \square

3.2. Operations. Let \mathcal{C} be an \mathcal{R} -stack. As for modules over a sheaf of algebras, there is a natural \mathcal{R} -functor

$$(3.2.1) \quad \otimes_{\mathcal{C}}: \text{Mod}(\mathcal{C}^{\text{op}}) \otimes_{\mathcal{R}} \text{Mod}(\mathcal{C}) \rightarrow \text{Mod}(\mathcal{R}).$$

This is discussed in [31] when X is reduced to a point. In order to explain how this extends to stacks, we need some preparation.

Denote by \mathcal{C}/X the Grothendieck construction associated with \mathcal{C} . Recall that objects of \mathcal{C}/X are pairs (u, γ) with $u: U \rightarrow X$ an open inclusion, and $\gamma \in \mathcal{C}(U)$. Morphisms $c: (u, \gamma) \rightarrow (u', \gamma')$ are defined only if $U' \subset U$, and in that case are given by morphisms $\gamma|_{U'} \rightarrow \gamma'$ in $\mathcal{C}(U')$. For $c': (u', \gamma') \rightarrow (u'', \gamma'')$ another morphism, the composition³ is given by $c' \circ c|_{U''}$.

³Here we denote for short by $c|_{U''}$ the composite $\gamma|_{U'} \xleftarrow{\sim} \gamma|_{U'|U''} \xrightarrow{c|_{U''}} \gamma'|_{U''}$.

Notation 3.2.1. Let \mathbf{C} be a category. We denote by $\text{Mor}_\circ \mathbf{C}$ the category whose objects are morphisms $c: \gamma \rightarrow \gamma'$ in \mathbf{C} and whose morphisms $c \rightarrow d$ are pairs (e, e') of morphisms in \mathbf{C} such that $c = e' \circ d \circ e$. This is visualised by the diagram

$$\begin{array}{ccc} \gamma & \xrightarrow{c} & \gamma' \\ e \downarrow & & \uparrow e' \\ \delta & \xrightarrow{d} & \delta' \end{array}$$

The following lemma is a stack-theoretical analogue of [22, Lemma 2.1.15].

Lemma 3.2.2. *Let $\mathcal{M}, \mathcal{N} \in \text{Mod}(\mathbf{C})$. Then*

(i) *the assignment*

$$((u, \gamma) \xrightarrow{c} (u', \gamma')) \mapsto \mathcal{H}om_{\mathcal{R}}(u_* \mathcal{M}(\gamma), u'_* \mathcal{N}(\gamma'))$$

defines a functor $(\text{Mor}_\circ \mathbf{C}/X)^{\text{op}} \rightarrow \text{Mod}(\mathcal{R})$;

(ii) *there is an isomorphism in $\text{Mod}(\mathcal{R})$*

$$(3.2.2) \quad \mathcal{H}om_{\text{Mod}(\mathbf{C})}(\mathcal{M}, \mathcal{N}) \simeq \varprojlim_{((u, \gamma) \xrightarrow{c} (u', \gamma')) \in \text{Mor}_\circ \mathbf{C}/X} \mathcal{H}om_{\mathcal{R}}(u_* \mathcal{M}(\gamma), u'_* \mathcal{N}(\gamma')).$$

Proof. (i) Let us check that a morphism (e, e') in $\text{Mor}_\circ \mathbf{C}/X$, visualised by the diagrams

$$\begin{array}{ccc} U & \longleftarrow & U' \\ \uparrow & & \downarrow \\ V & \longleftarrow & V' \end{array} \quad \begin{array}{ccc} (u, \gamma) & \xrightarrow{c} & (u', \gamma') \\ e \downarrow & & \uparrow e' \\ (v, \delta) & \xrightarrow{d} & (v', \delta') \end{array},$$

induces a morphism in $\text{Mod}(\mathcal{R})$

$$(3.2.3) \quad \mathcal{H}om_{\mathcal{R}}(v_* \mathcal{M}(\delta), v'_* \mathcal{N}(\delta')) \rightarrow \mathcal{H}om_{\mathcal{R}}(u_* \mathcal{M}(\gamma), u'_* \mathcal{N}(\gamma')).$$

Consider the inclusions of open subsets

$$\begin{array}{ccc} X & \xleftarrow{u} & U \\ & \swarrow v & \uparrow i \\ & & V. \end{array}$$

The morphism $e: \gamma|_V \rightarrow \delta$ induces a morphism

$$u_* \mathcal{M}(\gamma) \rightarrow u_* i_* i^{-1} \mathcal{M}(\gamma) \simeq v_* \mathcal{M}(\gamma|_V) \rightarrow v_* \mathcal{M}(\delta).$$

Similarly, $e': \delta'|_{U'} \rightarrow \gamma'$ induces a morphism $v'_* \mathcal{N}(\delta') \rightarrow u'_* \mathcal{N}(\gamma')$. Then (3.2.3) is obtained by left and right composition with these morphisms.

(ii) It is enough to prove that the natural morphism in $\text{Mod}(\mathcal{R}(X))$

$$\begin{aligned} \text{Hom}_{\text{Mod}(\mathcal{C})}(\mathcal{M}, \mathcal{N}) &\rightarrow \varprojlim_{c \in \text{Mor}_0 \mathcal{C}/X} \text{Hom}_{\mathcal{R}|_{U'}}(\mathcal{M}(\gamma|_{U'}), \mathcal{N}(\gamma')) \\ &\simeq \varprojlim_{c \in \text{Mor}_0 \mathcal{C}/X} \text{Hom}_{\mathcal{R}}(u_* \mathcal{M}(\gamma), u'_* \mathcal{N}(\gamma')) \end{aligned}$$

is an isomorphism. The proof of this fact follows the same arguments as those in the proof of [22, Lemma 2.1.15]. \square

For $\mathcal{M} \in \text{Mod}(\mathcal{C})$ and $\mathcal{P} \in \text{Mod}(\mathcal{C}^{\text{op}})$, similarly as above, the assignment

$$((u, \gamma) \xrightarrow{c} (u', \gamma')) \mapsto u'_! \mathcal{P}(\gamma') \otimes_{\mathcal{R}} u_* \mathcal{M}(\gamma)$$

defines a functor $\text{Mor}_0 \mathcal{C}/X \rightarrow \text{Mod}(\mathcal{R})$ and we set

$$(3.2.4) \quad \mathcal{P} \otimes_{\mathcal{C}} \mathcal{M} = \varinjlim_{((u, \gamma) \xrightarrow{c} (u', \gamma')) \in \text{Mor}_0 \mathcal{C}/X} u'_! \mathcal{P}(\gamma') \otimes_{\mathcal{R}} u_* \mathcal{M}(\gamma).$$

For \mathcal{A} an \mathcal{R} -algebra and $\mathcal{C} = \mathcal{A}^+$, this is the usual tensor product of $\mathcal{M} \in \text{Mod}(\mathcal{A})$ and $\mathcal{P} \in \text{Mod}(\mathcal{A}^{\text{op}})$. For example, when X is reduced to a point, this amounts to present $\mathcal{P} \otimes_{\mathcal{A}} \mathcal{M}$ as the quotient of $\bigoplus_{a \in \mathcal{A}} \mathcal{P} \otimes_{\mathcal{R}} \mathcal{M}$ by the subgroup generated by the elements $(p \otimes m)_a - (p \otimes am)_1$ and $(p \otimes m)_a - (pa \otimes m)_1$ for $a \in \mathcal{A}$, $p \in \mathcal{P}$ and $m \in \mathcal{M}$. (The indices denote the direct summand to which the elements belong. These generators correspond to the morphisms $(a, 1), (1, a): a \rightarrow 1$ in $\text{Mor}_0 \mathcal{A}^+$.)

Lemma 3.2.3. $\mathcal{P} \otimes_{\mathcal{C}} \mathcal{M}$ is a representative of the functor

$$\text{Hom}_{\text{Mod}(\mathcal{C})}(\mathcal{M}, \text{Hom}_{\mathcal{R}}(\mathcal{P}, \cdot)): \text{Mod}(\mathcal{R}) \rightarrow \text{Mod}(\mathcal{R}),$$

where, for any $\mathcal{L} \in \text{Mod}(\mathcal{R})$, we denote by $\text{Hom}_{\mathcal{R}}(\mathcal{P}, \mathcal{L})$ the \mathcal{C} -module $\text{Hom}_{\mathcal{R}}(\mathcal{P}(\cdot), \mathcal{L})$.

Proof. By Lemma 3.2.2, for any $\mathcal{L} \in \text{Mod}(\mathcal{R})$ there are isomorphisms

$$\begin{aligned} \text{Hom}_{\text{Mod}(\mathcal{C})}(\mathcal{M}, \text{Hom}_{\mathcal{R}}(\mathcal{P}, \mathcal{L})) &\simeq \varprojlim_{c \in \text{Mor}_0 \mathcal{C}/X} \text{Hom}_{\mathcal{R}}(u_* \mathcal{M}(\gamma), u'_! \text{Hom}_{\mathcal{R}|_{U'}}(\mathcal{P}(\gamma'), \mathcal{L}|_{U'})) \\ &\simeq \varprojlim_{c \in \text{Mor}_0 \mathcal{C}/X} \text{Hom}_{\mathcal{R}}(u_* \mathcal{M}(\gamma), \text{Hom}_{\mathcal{R}}(u'_! \mathcal{P}(\gamma'), \mathcal{L})) \\ &\simeq \varprojlim_{c \in \text{Mor}_0 \mathcal{C}/X} \text{Hom}_{\mathcal{R}}(u'_! \mathcal{P}(\gamma') \otimes_{\mathcal{R}} u_* \mathcal{M}(\gamma), \mathcal{L}) \\ &\simeq \text{Hom}_{\mathcal{R}} \left(\varinjlim_{c \in \text{Mor}_0 \mathcal{C}/X} u'_! \mathcal{P}(\gamma') \otimes_{\mathcal{R}} u_* \mathcal{M}(\gamma), \mathcal{L} \right) \\ &= \text{Hom}_{\mathcal{R}}(\mathcal{P} \otimes_{\mathcal{C}} \mathcal{M}, \mathcal{L}). \end{aligned}$$

\square

Uniqueness of representatives imply the functoriality of the assignment $(\mathcal{P}, \mathcal{M}) \mapsto \mathcal{P} \otimes_{\mathcal{C}} \mathcal{M}$. We have thus defined the functor (3.2.1).

As for modules over a sheaf of algebras, let us use the short hand notation

$$(3.2.5) \quad \mathcal{H}om_{\mathcal{C}}(\cdot, \cdot): \text{Mod}(\mathcal{C})^{\text{op}} \otimes_{\mathcal{R}} \text{Mod}(\mathcal{C}) \rightarrow \text{Mod}(\mathcal{R}).$$

for the internal hom-functor $\mathcal{H}om_{\text{Mod}(\mathcal{C})}(\cdot, \cdot)$.

Notation 3.2.4. We use the same notation

$$\begin{aligned} \mathcal{H}om_{\mathcal{C}}: \text{Mod}(\mathcal{C} \otimes_{\mathcal{R}} \mathcal{D}^{\text{op}})^{\text{op}} \otimes_{\mathcal{R}} \text{Mod}(\mathcal{C} \otimes_{\mathcal{R}} \mathcal{E}) &\rightarrow \text{Mod}(\mathcal{D} \otimes_{\mathcal{R}} \mathcal{E}), \\ \otimes_{\mathcal{C}}: \text{Mod}(\mathcal{C}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{D}) \otimes_{\mathcal{R}} \text{Mod}(\mathcal{C} \otimes_{\mathcal{R}} \mathcal{E}) &\rightarrow \text{Mod}(\mathcal{D} \otimes_{\mathcal{R}} \mathcal{E}) \end{aligned}$$

for the \mathcal{R} -functors obtained by “picking up operators” (in the sense of [30, page 15]) from the \mathcal{R} -functors (3.2.5) and (3.2.1).

These functors satisfy the relations (3.2.2) and (3.2.4). For example, let $\mathcal{P} \in \text{Mod}(\mathcal{C}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{D})$ and $\mathcal{M} \in \text{Mod}(\mathcal{C} \otimes_{\mathcal{R}} \mathcal{E})$. Consider them as functors $\mathcal{P}: \mathcal{C}^{\text{op}} \rightarrow \text{Mod}(\mathcal{D})$ and $\mathcal{M}: \mathcal{C} \rightarrow \text{Mod}(\mathcal{E})$. Then $\mathcal{P} \otimes_{\mathcal{C}} \mathcal{M}$ satisfies (3.2.4), where the operations $u'_!$ and $u_!$ are those of the proper stacks $\text{Mod}(\mathcal{D})$ and $\text{Mod}(\mathcal{E})$, respectively, and the tensor product is the natural functor

$$\otimes_{\mathcal{R}}: \text{Mod}(\mathcal{D}) \otimes_{\mathcal{R}} \text{Mod}(\mathcal{E}) \rightarrow \text{Mod}(\mathcal{D} \otimes_{\mathcal{R}} \mathcal{E}).$$

The standard formulas concerning the usual hom-functor and tensor product hold. For example,

Lemma 3.2.5. *For $\mathcal{M} \in \text{Mod}(\mathcal{C}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{D})$, $\mathcal{N} \in \text{Mod}(\mathcal{C} \otimes_{\mathcal{R}} \mathcal{E})$, and $\mathcal{P} \in \text{Mod}(\mathcal{D} \otimes_{\mathcal{R}} \mathcal{F})$, there is an isomorphism in $\text{Mod}(\mathcal{E}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{F})$*

$$\mathcal{H}om_{\mathcal{D}}(\mathcal{M} \otimes_{\mathcal{C}} \mathcal{N}, \mathcal{P}) \simeq \mathcal{H}om_{\mathcal{C}}(\mathcal{N}, \mathcal{H}om_{\mathcal{D}}(\mathcal{M}, \mathcal{P})).$$

Recall the notation $\mathcal{C} \in \text{Mod}(\mathcal{C}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{C})$ in (3.1.2). By Lemma 3.1.3 the functors $\mathcal{H}om_{\mathcal{C}}(\mathcal{C}, \cdot)$, and hence also $\mathcal{C} \otimes_{\mathcal{C}} (\cdot)$, are isomorphic to the identity.

Using (3.2.4) with $\mathcal{P} = \mathcal{C}$, we get

Lemma 3.2.6. *For $\mathcal{M} \in \text{Mod}(\mathcal{C})$ there is an isomorphism in $\text{Mod}(\mathcal{C})$*

$$\mathcal{M} \simeq \varinjlim_{((u, \gamma) \xrightarrow{c} (u', \gamma')) \in \text{Mor}_o \mathcal{C}/X} u'_! \mathcal{Y}_{\mathcal{C}}(\gamma') \otimes_{\mathcal{R}} u_* \mathcal{M}(\gamma).$$

3.3. Morita equivalence. Let us discuss how classical Morita theory extends to linear stacks.

Let \mathcal{C} and \mathcal{D} be \mathcal{R} -stacks. Denote by $\text{Fct}_{\mathcal{R}}^r(\text{Mod}(\mathcal{C}), \text{Mod}(\mathcal{D}))$ the stack of \mathcal{R} -functors admitting a right adjoint. The Eilenberg-Watts theorem for \mathcal{R} -stacks holds:

Theorem 3.3.1. (i) *The \mathcal{R} -functor*

$$\text{Mod}(\mathcal{C}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{D}) \rightarrow \text{Fct}_{\mathcal{R}}^r(\text{Mod}(\mathcal{C}), \text{Mod}(\mathcal{D})), \quad \mathcal{P} \mapsto \mathcal{P} \otimes_{\mathcal{C}} (\cdot),$$

is an equivalence.

- (ii) For $\mathcal{P} \in \text{Mod}(\mathcal{C}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{D})$ and $\mathcal{Q} \in \text{Mod}(\mathcal{D}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{E})$ one has an isomorphism in $\text{Fct}_{\mathcal{R}}^r(\text{Mod}(\mathcal{C}), \text{Mod}(\mathcal{E}))$

$$(\mathcal{Q} \otimes_{\mathcal{D}} \mathcal{P}) \otimes_{\mathcal{C}} (\cdot) \simeq (\mathcal{Q} \otimes_{\mathcal{D}} (\cdot)) \circ (\mathcal{P} \otimes_{\mathcal{C}} (\cdot)).$$

Given an \mathcal{R} -functor $\mathbf{h}: \text{Mod}(\mathcal{C}) \rightarrow \text{Mod}(\mathcal{D})$, we will use the same notation \mathbf{h} for the induced \mathcal{R} -functor, obtained by “picking up operators”,

$$\text{Mod}(\mathcal{C}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{C}) \rightarrow \text{Mod}(\mathcal{C}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{D}).$$

Proof. (i) Let us show that $\mathbf{h} \mapsto \mathbf{h}(\mathcal{C})$ is a quasi-inverse to the functor in the statement.

Since $\mathcal{P} \simeq \mathcal{P} \otimes_{\mathcal{C}} \mathcal{C}$, we only have to prove that $\mathbf{h} \xrightarrow{\sim} \mathbf{h}(\mathcal{C}) \otimes_{\mathcal{C}} (\cdot)$. Let $\mathcal{M} \in \text{Mod}(\mathcal{C})$. Since \mathbf{h} has a right adjoint, it commutes with direct limits, proper direct images, and tensor products with \mathcal{R} -modules. Hence we have

$$\begin{aligned} \mathbf{h}(\mathcal{M}) &\simeq \mathbf{h} \left(\varinjlim_{c \in \text{Mor}_o \mathcal{C}/X} u'_! Y_{\mathcal{C}}(\gamma') \otimes_{\mathcal{R}} u_* \mathcal{M}(\gamma) \right) \\ &\simeq \varinjlim_{c \in \text{Mor}_o \mathcal{C}/X} u'_! \mathbf{h}(Y_{\mathcal{C}}(\gamma')) \otimes_{\mathcal{R}} u_* \mathcal{M}(\gamma) \\ &\simeq \varinjlim_{c \in \text{Mor}_o \mathcal{C}/X} u'_! \mathbf{h}(\mathcal{C})(\gamma', \cdot) \otimes_{\mathcal{R}} u_* \mathcal{M}(\gamma) \\ &= \mathbf{h}(\mathcal{C}) \otimes_{\mathcal{C}} \mathcal{M}, \end{aligned}$$

where the first isomorphism follows from Lemma 3.2.6 and we use the fact that $\mathbf{h} \circ Y_{\mathcal{C}}$ identifies to $\mathbf{h}(\mathcal{C})$ via the equivalence induced by Lemma 2.1.1.

- (ii) By (i), the statement amounts to the isomorphism in $\text{Mod}(\mathcal{C}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{E})$

$$(\mathcal{Q} \otimes_{\mathcal{D}} \mathcal{P}) \otimes_{\mathcal{C}} \mathcal{C} \simeq \mathcal{Q} \otimes_{\mathcal{D}} (\mathcal{P} \otimes_{\mathcal{C}} \mathcal{C}).$$

□

Remark 3.3.2. Denoting by $\text{Fct}_{\mathcal{R}}^l(\text{Mod}(\mathcal{C}), \text{Mod}(\mathcal{D}))$ the stack of \mathcal{R} -functors admitting a left adjoint, one similarly gets an \mathcal{R} -equivalence

$$\text{Mod}(\mathcal{C}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{D}) \rightarrow \text{Fct}_{\mathcal{R}}^l(\text{Mod}(\mathcal{D}), \text{Mod}(\mathcal{C}))^{\text{op}}, \quad \mathcal{P} \mapsto \mathcal{H}om_{\mathcal{D}}(\mathcal{P}, \cdot),$$

and the corresponding commutative diagram as in Theorem 3.3.1 (ii). These constructions are interchanged by the \mathcal{R} -equivalence

$$\text{Fct}_{\mathcal{R}}^r(\text{Mod}(\mathcal{C}), \text{Mod}(\mathcal{D})) \approx_{\mathcal{R}} \text{Fct}_{\mathcal{R}}^l(\text{Mod}(\mathcal{D}), \text{Mod}(\mathcal{C}))^{\text{op}}$$

sending a functor to its adjoint.

Definition 3.3.3. (i) One says that $\mathcal{Q} \in \text{Mod}(\mathcal{D}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{C})$ is an inverse of $\mathcal{P} \in \text{Mod}(\mathcal{C}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{D})$ if there are isomorphisms of $\mathcal{C} \otimes_{\mathcal{R}} \mathcal{C}^{\text{op}}$ - and $\mathcal{D} \otimes_{\mathcal{R}} \mathcal{D}^{\text{op}}$ -modules, respectively,

$$\mathcal{Q} \otimes_{\mathcal{D}} \mathcal{P} \simeq \mathcal{C}, \quad \mathcal{P} \otimes_{\mathcal{C}} \mathcal{Q} \simeq \mathcal{D}.$$

- (ii) An object $\mathcal{P} \in \text{Mod}(\mathcal{C}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{D})$ is called invertible if it has an inverse.

One proves (see e.g. [22, §19.5]) that \mathcal{P} is invertible if and only if one of the following equivalent conditions is satisfied

- (i) $\mathcal{H}om_{\mathbf{D}}(\mathcal{P}, \mathbf{D})$ is an inverse of \mathcal{P} ;
- (ii) the functor $\mathcal{P} \otimes_{\mathbf{C}} (\cdot): \mathbf{Mod}(\mathbf{C}) \rightarrow \mathbf{Mod}(\mathbf{D})$ is an \mathcal{R} -equivalence.
- (iii) the functor $\mathcal{H}om_{\mathbf{C}^{\text{op}}}(\mathcal{P}, \cdot): \mathbf{Mod}(\mathbf{C}^{\text{op}}) \rightarrow \mathbf{Mod}(\mathbf{D}^{\text{op}})$ is an \mathcal{R} -equivalence.

Notation 3.3.4. For any \mathcal{R} -functor $\mathbf{f}: \mathbf{C} \rightarrow \mathbf{C}'$, we denote by $\mathbf{End}_{\mathbf{C}'}(\mathbf{f})$ the \mathcal{R} -stack associated to the separated prestack whose objects on $U \subset X$ are those of $\mathbf{C}(U)$ and $\mathbf{Hom}(\gamma, \gamma') = \mathbf{Hom}_{\mathbf{C}'(U)}(\mathbf{f}(\gamma), \mathbf{f}(\gamma'))$.

Note that, if \mathbf{f} is fully faithful, then the natural \mathcal{R} -functor $\mathbf{C} \rightarrow \mathbf{End}_{\mathbf{C}'}(\mathbf{f})$ induced by \mathbf{f} is an equivalence. In particular, identifying $\mathbf{C} \in \mathbf{Mod}(\mathbf{C}^{\text{op}} \otimes_{\mathcal{R}} \mathbf{C})$ with the Yoneda embedding $\mathbf{Y}_{\mathbf{C}}: \mathbf{C}^{\text{op}} \rightarrow \mathbf{Mod}(\mathbf{C})$, one has $\mathbf{C}^{\text{op}} \approx_{\mathcal{R}} \mathbf{End}_{\mathbf{Mod}(\mathbf{C})}(\mathbf{C})$. Moreover, considering $\mathcal{P} \in \mathbf{Mod}(\mathbf{C}^{\text{op}} \otimes_{\mathcal{R}} \mathbf{D})$ as a functor $\mathbf{C}^{\text{op}} \rightarrow \mathbf{Mod}(\mathbf{D})$, the condition of \mathcal{P} being invertible is further equivalent to

- (iv) \mathcal{P} is a faithfully flat⁴ \mathbf{D} -module locally of finite presentation⁵ and $\mathbf{C}^{\text{op}} \approx_{\mathcal{R}} \mathbf{End}_{\mathbf{Mod}(\mathbf{D})}(\mathcal{P})$;
- (v) \mathcal{P} is \mathbf{D} -progenerator⁶ locally of finite type and $\mathbf{C}^{\text{op}} \approx_{\mathcal{R}} \mathbf{End}_{\mathbf{Mod}(\mathbf{D})}(\mathcal{P})$.

By reversing the role of \mathbf{C}^{op} and \mathbf{D} , one gets dual equivalent conditions.

Theorem 3.3.5 (Morita). *An \mathcal{R} -functor $\mathbf{h}: \mathbf{Mod}(\mathbf{C}) \rightarrow \mathbf{Mod}(\mathbf{D})$ is an equivalence if and only if $\mathcal{P} = \mathbf{h}(\mathbf{C})$ is an invertible $(\mathbf{C}^{\text{op}} \otimes_{\mathcal{R}} \mathbf{D})$ -module. Moreover, one has $\mathbf{h} \simeq \mathcal{P} \otimes_{\mathbf{C}} (\cdot)$.*

Definition 3.3.6. Two \mathcal{R} -stacks \mathbf{C} and \mathbf{D} are Morita \mathcal{R} -equivalent if their stacks of modules $\mathbf{Mod}(\mathbf{C})$ and $\mathbf{Mod}(\mathbf{D})$ are \mathcal{R} -equivalent.

Hence \mathbf{C} and \mathbf{D} are Morita \mathcal{R} -equivalent if and only if there exists an invertible $(\mathbf{C}^{\text{op}} \otimes_{\mathcal{R}} \mathbf{D})$ -module.

Let us say that $\mathcal{P} \in \mathbf{Mod}(\mathbf{C}^{\text{op}} \otimes_{\mathcal{R}} \mathbf{D}) \approx_{\mathcal{R}} \mathbf{Fct}_{\mathcal{R}}(\mathbf{D}, \mathbf{Mod}(\mathbf{C}^{\text{op}}))$ is locally free of rank one over \mathbf{C}^{op} if for any $\delta \in \mathbf{D}$ the \mathbf{C}^{op} -module $\mathcal{P}(\delta)$ is locally free of rank one, that is to say, the functor $\mathcal{P}(\delta): \mathbf{C}^{\text{op}} \rightarrow \mathbf{Mod}(\mathcal{R})$ is representable.

Recall from (2.2.2) that $_{\mathbf{f}}(\cdot): \mathbf{Mod}(\mathbf{C}) \rightarrow \mathbf{Mod}(\mathbf{D})$ denotes the functor associated to an \mathcal{R} -functor $\mathbf{f}: \mathbf{D} \rightarrow \mathbf{C}$.

Proposition 3.3.7. *The \mathcal{R} -functor*

$$(3.3.1) \quad \mathbf{Fct}_{\mathcal{R}}(\mathbf{D}, \mathbf{C}) \rightarrow \mathbf{Mod}(\mathbf{C}^{\text{op}} \otimes_{\mathcal{R}} \mathbf{D}), \quad \mathbf{f} \mapsto _{\mathbf{f}}\mathbf{C}$$

is fully faithful and induces an equivalence with the full substack of locally free modules of rank one over \mathbf{C}^{op} .

⁴ \mathcal{P} is a faithfully flat \mathbf{D} -module if the functor $(\cdot) \otimes_{\mathbf{D}} \mathcal{P}$ is faithful and exact.

⁵ \mathcal{P} is a \mathbf{D} -module of finite presentation if the functor $\mathcal{H}om_{\mathbf{D}}(\mathcal{P}, \cdot)$ commutes with small filtrant colimits.

⁶ \mathcal{P} is \mathbf{D} -progenerator if the functor $\mathcal{H}om_{\mathbf{D}}(\mathcal{P}, \cdot)$ is faithful and exact.

Proof. (i) The functor in the statement equals $Y_{C^{\text{op}}} \circ \cdot$. This is fully faithful, since $Y_{C^{\text{op}}}$ is fully faithful.

(ii) Assume that $\mathcal{P} \in \text{Mod}(C^{\text{op}} \otimes_{\mathcal{R}} D)$ is a locally free module of rank one over C^{op} . Then $\mathcal{P} \simeq {}_f C$, where $f: D \rightarrow C$ is the functor associating to $\delta \in D$ the representative of $\mathcal{P}(\delta)$. \square

Corollary 3.3.8. *Two \mathcal{R} -stacks C and D are \mathcal{R} -equivalent if and only if there exists $\mathcal{P} \in \text{Mod}(C^{\text{op}} \otimes_{\mathcal{R}} D)$ which is invertible and locally free of rank one over C^{op} .*

In particular, two algebroids A and B are \mathcal{R} -equivalent if and only if there exists an invertible $(A^{\text{op}} \otimes_{\mathcal{R}} B)$ -module \mathcal{P} which is locally free of rank one over A^{op} . These conditions on \mathcal{P} are equivalent to the condition that \mathcal{P} is bi-invertible in the sense of [23, Corollary 2.1.10].

Remark 3.3.9. If $C \approx_{\mathcal{R}} \mathcal{A}^+$ and $D \approx_{\mathcal{R}} \mathcal{B}^+$, the functor $\mathcal{B}^+ \rightarrow \mathcal{A}^+$ associated to an $\mathcal{A}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{B}$ -module \mathcal{P} locally free of rank one over \mathcal{A}^{op} is $f = (\cdot) \otimes_{\mathcal{B}} \mathcal{P}$. Note that any local isomorphism $h: \mathcal{A} \xrightarrow{\sim} \mathcal{P}$ of right \mathcal{A} -modules defines a local \mathcal{R} -algebra morphism (isomorphism if \mathcal{P} invertible)

$$(3.3.2) \quad f: \mathcal{B} \rightarrow \text{End}_{\mathcal{A}^{\text{op}}}(\mathcal{P}) \xrightarrow{\text{Ad}(h^{-1})} \text{End}_{\mathcal{A}^{\text{op}}}(\mathcal{A}) \simeq \mathcal{A},$$

(the first arrow is induced by the \mathcal{B} -module structure of \mathcal{P}), for which $h: {}_f \mathcal{A} \xrightarrow{\sim} \mathcal{P}$ is an isomorphism of $\mathcal{A}^{\text{op}} \otimes_{\mathcal{R}} \mathcal{B}$ -modules and $f \simeq f^+$. If h is given by $a \mapsto ua$ for a local generator u of the right \mathcal{A} -modules \mathcal{P} , then $f(b) = a$ for a such that $ua = bu$.

3.4. Picard good stacks. We will use the notation

$$C^e = C^{\text{op}} \otimes_{\mathcal{R}} C.$$

Definition 3.4.1. An \mathcal{R} -stack C is Picard good if all invertible C^e -modules are locally free of rank one over C^{op} (or, equivalently, over C). An \mathcal{R} -algebra \mathcal{A} is Picard good if it is so as an \mathcal{R} -stack.

For $\mathcal{A} = \mathcal{R}$, one recovers Definition 4.2 given in [11].

Since the condition of being Picard good is local, an algebroid is Picard good if and only if so are the algebras that locally represent it.

Recall from (v) in Section 3.3 that invertible bimodules are projective as right (or left) modules. It follows that examples of Picard good rings are projective-free rings, and in particular local rings. Note however that Picard good does not imply projective-free (see Remark 4.3.3).

Denote by $\text{Inv}(C^e)$ the substack of $\text{Mod}(C^e)$ whose objects are invertible C^e -modules and whose morphisms are only those morphisms which are invertible. Then \otimes_C induces on $\text{Inv}(C^e)$ a natural structure of stack of 2-groups, and (3.3.1) gives a fully faithful functor of stacks of 2-groups

$$(3.4.1) \quad \text{Aut}_{\mathcal{R}}(C)_{\text{op}} \hookrightarrow \text{Inv}(C^e), \quad f \mapsto {}_f C.$$

Here, for \mathbf{G} a stack of 2-groups, \mathbf{G}_{op} denotes the stack of 2-groups with the same groupoid structure as \mathbf{G} and with reversed monoidal structure. Set

$$\mathcal{O}ut_{\mathcal{R}}(\mathbf{C}) = \pi_0(\mathbf{Aut}_{\mathcal{R}}(\mathbf{C})), \quad \mathcal{P}ic_{\mathcal{R}}(\mathbf{C}) = \pi_0(\mathbf{Inv}(\mathbf{C}^e)).$$

Then (3.4.1) induces an injective homomorphism of groups

$$(3.4.2) \quad \mathcal{O}ut_{\mathcal{R}}(\mathbf{C})^{\text{op}} \hookrightarrow \mathcal{P}ic_{\mathcal{R}}(\mathbf{C}).$$

Note that, from (2.5.1) follows that $\mathcal{O}ut(\mathcal{A}^+) = \mathcal{A}ut_{\mathcal{R}}(\mathcal{A})/\mathcal{I}nn(\mathcal{A})$, the sheaf of outer automorphisms of \mathcal{A} .

Proposition 3.4.2. *Let \mathbf{C} be an \mathcal{R} -stack. Then the following are equivalent*

- (i) \mathbf{C} is Picard good;
- (ii) (3.4.1) is an equivalence;
- (iii) (3.4.2) is an isomorphism.

Proof. The equivalence between (i) and (ii) follows from Proposition 3.3.7, whereas that between (ii) and (iii) follows from the fact that a functor of stacks $\mathbf{C} \rightarrow \mathbf{C}'$ is essentially surjective if and only if the induced morphism of sheaves $\pi_0(\mathbf{C}) \rightarrow \pi_0(\mathbf{C}')$ is surjective. \square

Proposition 3.4.3. *Let \mathbf{C} be a Picard good \mathcal{R} -stack.*

- (i) *Let \mathbf{D} be an \mathcal{R} -stack locally equivalent to \mathbf{C} . Then \mathbf{C} and \mathbf{D} are Morita \mathcal{R} -equivalent if and only if they are \mathcal{R} -equivalent.*
- (ii) *Let \mathbf{M} be an \mathcal{R} -stack locally \mathcal{R} -equivalent to $\mathbf{Mod}(\mathbf{C})$. Then $\mathbf{M} \approx_{\mathcal{R}} \mathbf{Mod}(\mathbf{D})$ for an \mathcal{R} -stack \mathbf{D} locally \mathcal{R} -equivalent to \mathbf{C} .*

Proof. (i) By Theorem 3.3.5, there is an equivalence of stacks of 2-groups

$$\mathbf{Inv}(\mathbf{C}^e) \xrightarrow{\sim} \mathbf{Aut}_{\mathcal{R}}(\mathbf{Mod}(\mathbf{C})), \quad \mathcal{P} \mapsto \mathcal{P} \otimes_{\mathbf{C}} (\cdot).$$

We thus have a (quasi-)commutative diagram

$$\begin{array}{ccc} \mathbf{Inv}(\mathbf{C}^e) & \xrightarrow{\sim} & \mathbf{Aut}_{\mathcal{R}}(\mathbf{Mod}(\mathbf{C})) \\ & \searrow & \nearrow \mathbf{m} \\ & \mathbf{Aut}_{\mathcal{R}}(\mathbf{C})_{\text{op}} & \end{array}$$

where \mathbf{m} is induced by the functor $\mathbf{Mod}(\cdot)$. It follows from Proposition 3.4.2 that \mathbf{C} is Picard good if and only if \mathbf{m} is an equivalence.

Let $\mathbf{Equiv}_{\mathcal{R}}(\cdot, \cdot)$ denote the stack of \mathcal{R} -equivalences, with invertible transformations as morphisms. Consider the functor

$$\mathbf{Equiv}_{\mathcal{R}}(\mathbf{C}, \mathbf{D}) \rightarrow \mathbf{Equiv}_{\mathcal{R}}(\mathbf{Mod}(\mathbf{D}), \mathbf{Mod}(\mathbf{C}))$$

induced by the 2-functor $\mathbf{Mod}(\cdot)$. Since \mathbf{D} is locally equivalent to \mathbf{C} , this locally reduces to the functor \mathbf{m} above. It is thus locally, hence globally, an equivalence.

(ii) Let $\mathbf{E} \subset \mathbf{M}$ be the full substack of objects γ with the property that for any local \mathcal{R} -equivalence $\mathbf{h}: \mathbf{M} \xrightarrow{\sim} \mathbf{Mod}(\mathbf{C})$, the \mathbf{C} -module $\mathbf{h}(\gamma)$ is locally free of rank one. Since \mathbf{C} is Picard good, the \mathcal{R} -stack \mathbf{E} is locally non-empty and locally \mathcal{R} -equivalent to \mathbf{C}^{op} . Set $\mathbf{D} = \mathbf{E}^{\text{op}}$. Then the \mathcal{R} -functor

$$\mathbf{M} \rightarrow \mathbf{Mod}(\mathbf{D}), \quad \delta \mapsto \mathcal{H}om_{\mathbf{M}}(\cdot, \delta)$$

is locally, hence globally, an equivalence. \square

If \mathbf{C} is an invertible \mathcal{R} -algebroid, then it is Picard good if and only if \mathcal{R} is, and one has equivalences of stacks of 2-groups

$$(3.4.3) \quad \mathcal{R}^{\times}[1] \xrightarrow{\sim} \text{Inv}(\mathcal{R}) \approx \text{Inv}(\mathbf{C}^e), \quad \mathcal{P} \mapsto \mathcal{R} \times_{\mathcal{R}^{\times}} \mathcal{P}.$$

(Recall that $\mathcal{R}^{\times}[1]$ denotes the stack of \mathcal{R}^{\times} -torsors.) Moreover, in this situation the stack \mathbf{D} in (ii) above is \mathcal{R} -equivalent via $\gamma \mapsto \mathcal{H}om_{\mathbf{M}}(\gamma, \cdot)$ to the full substack of $\mathbf{Fct}_{\mathcal{R}}(\mathbf{M}, \mathbf{Mod}(\mathcal{R}))$ whose objects are equivalences.

Examples of stacks as in Proposition 3.4.3 (ii) arise from deformations of categories of modules as discussed in [28]. In particular, Proposition 3.4.3 applies when \mathbf{C} is (equivalent to) the structure sheaf of a ringed space. We thus recover results of [27].

4. MICRODIFFERENTIAL OPERATORS

We collect here some results from the theory of microdifferential operators of [37] (see also [38, 16, 18]). The statements about the automorphisms of the sheaf of microdifferential operators are well known. Since we lack a reference for the proofs, we give them here.

4.1. Microdifferential operators. Let M be an n -dimensional complex manifold, T^*M its cotangent bundle and $\dot{T}^*M \subset T^*M$ the open subset obtained by removing the zero-section.

Denote by $\mathcal{E}_{\dot{T}^*M}$ the sheaf of microdifferential operators on \dot{T}^*M . Recall that $\mathcal{E}_{\dot{T}^*M}$ is a sheaf of central \mathbb{C} -algebras endowed with a \mathbb{Z} -filtration by the order of the operators, and one has

$$\mathcal{G}r \mathcal{E}_{\dot{T}^*M} \simeq \bigoplus_{m \in \mathbb{Z}} \mathcal{O}_{\dot{T}^*M}(m),$$

where $\mathcal{O}_{\dot{T}^*M}(m)$ is the subsheaf of $\mathcal{O}_{\dot{T}^*M}$ of holomorphic functions homogeneous of degree m .

For $\lambda \in \mathbb{C}$, denote by $\mathcal{E}_{\dot{T}^*M}(\lambda)$ the sheaf of microdifferential operators of order at most λ . In a local coordinate system (x) on M , with associated symplectic coordinates $(x; \xi)$ on \dot{T}^*M , a section $P \in \Gamma(V; \mathcal{E}_{\dot{T}^*M}(\lambda))$ is determined by its total symbol, which is a formal series

$$\text{tot}(P) = \sum_{j=0}^{+\infty} p_{\lambda-j}(x, \xi)$$

with $p_{\lambda-j} \in \Gamma(V; \mathcal{O}_{\dot{T}^*M})$ homogeneous of degree $\lambda - j$, satisfying suitable growth conditions in j . If Q is a section of $\mathcal{E}_{\dot{T}^*M}(\mu)$, then $PQ \in \mathcal{E}_{\dot{T}^*M}(\lambda + \mu)$ has total symbol given by the Leibniz formula

$$\text{tot}(PQ) = \sum_{\alpha \in \mathbb{N}^n} \frac{1}{\alpha!} \partial_\xi^\alpha \text{tot}(P) \partial_x^\alpha \text{tot}(Q).$$

Set

$$\mathcal{E}_{\dot{T}^*M}^{[\lambda]} = \bigcup_{n \in \mathbb{Z}} \mathcal{E}_{\dot{T}^*M}(\lambda + n),$$

where $[\lambda]$ is the class of λ in \mathbb{C}/\mathbb{Z} , and denote by

$$\sigma_\lambda: \mathcal{E}_{\dot{T}^*M}(\lambda) \rightarrow \mathcal{O}_{\dot{T}^*M}(\lambda) \quad \text{and} \quad \sigma: \mathcal{E}_{\dot{T}^*M}^{[\lambda]} \rightarrow \mathcal{O}_{\dot{T}^*M}$$

the symbol of order λ and the principal symbol, respectively, where $\sigma(P) = \sigma_\mu(P)$ for $P \in \mathcal{E}_{\dot{T}^*M}(\mu) \setminus \mathcal{E}_{\dot{T}^*M}(\mu - 1)$. Note that $\mathcal{E}_{\dot{T}^*M}^{[\lambda]}$ is a bimodule over $\mathcal{E}_{\dot{T}^*M} = \mathcal{E}_{\dot{T}^*M}^{[0]}$ and for any $P \in \mathcal{E}_{\dot{T}^*M}(\lambda)$ and $Q \in \mathcal{E}_{\dot{T}^*M}(\mu)$ one has

$$\sigma_{\lambda+\mu}(PQ) = \sigma_\lambda(P) \sigma_\mu(Q).$$

Recall that a microdifferential operator is invertible at $p \in \dot{T}^*M$ if and only if its principal symbol does not vanish at p .

4.2. Automorphisms of $\mathcal{E}_{\dot{T}^*M}$.

Lemma 4.2.1. *Any \mathbb{C} -algebra automorphism of $\mathcal{E}_{\dot{T}^*M}$ is filtered and symbol preserving.*

Proof. Let f be a \mathbb{C} -algebra automorphism of $\mathcal{E}_{\dot{T}^*M}$. Define the spectrum of $P \in \Gamma(V; \mathcal{E}_{\dot{T}^*M})$ as

$$\Sigma(P): V \rightarrow \mathcal{P}(\mathbb{C})$$

$$p \mapsto \{a \in \mathbb{C}: a - P \text{ is not invertible at } p\},$$

where $\mathcal{P}(\mathbb{C})$ denotes the set of subsets of \mathbb{C} . Note that $\Sigma(P) = \Sigma(f(P))$. Set for short

$$\mathcal{E}_m = \mathcal{E}_{\dot{T}^*M}(m) \setminus \mathcal{E}_{\dot{T}^*M}(m - 1).$$

Recall that P is invertible if and only if its principal symbol does not vanish.

(i) If $P \in \mathcal{E}_0$ and its principal symbol is not locally constant, then $\Sigma(P)(p) = \{\sigma(P)(p)\}$. Since $\Sigma(P) = \Sigma(f(P))$, it follows that $f(P) \in \mathcal{E}_0$ and $\sigma(P) = \sigma(f(P))$.

(ii) Let $P \in \mathcal{E}_0$ have locally constant principal symbol. For any $Q \in \mathcal{E}_{\dot{T}^*M}(0) \setminus \sigma_0^{-1}(\mathbb{C}_{\dot{T}^*M})$ one has

$$\sigma_0(P) \sigma_0(Q) = \sigma_0(PQ) = \sigma_0(f(PQ)) = \sigma(f(P)) \sigma_0(f(Q)) = \sigma(f(P)) \sigma_0(Q)$$

where the second equality follows from (i). One deduces $\sigma(f(P)) = \sigma_0(P)$, so that in particular $f(P) \in \mathcal{E}_0$.

(iii) Pick an operator $D \in \mathcal{E}_1$ invertible at p , and let d be the order of $f(D)$. Then $f(D)^m$ is an invertible operator of order dm and one has

$$f(\mathcal{E}_m) = f(D^m \mathcal{E}_0) = f(D)^m f(\mathcal{E}_0) = f(D)^m \mathcal{E}_0 = \mathcal{E}_{dm}.$$

Since f is an automorphism of $\mathcal{E}_{\dot{T}^*M} \setminus \{0\} = \bigsqcup_{m \in \mathbb{Z}} \mathcal{E}_m$, it follows that $d = \pm 1$. Thus f either preserves or reverses the order. Note that if an operator P satisfies $\sigma(P)(p) = 0$, then $\Sigma(P)(p) = \mathbb{C}$ if and only if P has positive order. Hence, f preserves the order.

(iv) We have proved that f is filtered and preserves the symbol of operators in \mathcal{E}_0 . As $\mathcal{E}_m = D^m \mathcal{E}_0$, to show that f is symbol preserving it is enough to check that $\sigma_1(D) = \sigma_1(f(D))$.

Let $(x; \xi)$ be a local system of symplectic coordinates at p . Identifying x_i with the operator in \mathcal{E}_0 whose total symbol is x_i , one has

$$\begin{aligned} \partial_{\xi_i} \sigma_1(D) &= \{x_i, \sigma_1(D)\} = \{\sigma_0(x_i), \sigma_1(D)\} = \sigma_0([x_i, D]) \\ &= \sigma_0(f([x_i, D])) = \sigma_0([f(x_i), f(D)]) = \{\sigma_0(f(x_i)), \sigma_1(f(D))\} \\ &= \{x_i, \sigma_1(f(D))\} = \partial_{\xi_i} \sigma_1(f(D)), \quad \text{for } i = 1, \dots, n, \end{aligned}$$

so that

$$\sigma_1(D) = \sigma_1(f(D)) + \varphi(x),$$

and one takes the homogeneous component of degree 1. \square

Proposition 4.2.2. *Any \mathbb{C} -algebra automorphism of $\mathcal{E}_{\dot{T}^*M}$ is locally of the form $\text{Ad}(P)$ for some $\lambda \in \mathbb{C}$ and some invertible $P \in \mathcal{E}_{\dot{T}^*M}(\lambda)$.*

Proof. Identify $\dot{T}^*M \times \dot{T}^*M$ to an open subset of $T^*(M \times M)$. Let (x) be a system of local coordinates on M , and denote by (x, y) the coordinates on $M \times M$. For $Q \in \mathcal{E}_{\dot{T}^*M}$, denote by Q_x and Q_y its pull-back to $\mathcal{E}_{\dot{T}^*M \times \dot{T}^*M}$ by the first and second projection, respectively.

Let $f: \mathcal{E}_{\dot{T}^*M} \rightarrow \mathcal{E}_{\dot{T}^*M}$ be a \mathbb{C} -algebra automorphism. By Lemma 4.2.1, f is filtered and symbol preserving. Denote by \mathcal{L} the $\mathcal{E}_{\dot{T}^*M \times \dot{T}^*M}$ -module with one generator u and relations

$$(x_i - f(y_i))u = (\partial_{x_i} - f(\partial_{y_i}))u = 0, \quad \text{for } i = 1, \dots, n.$$

Then the image $f(Q)$ of $Q \in \mathcal{E}_{\dot{T}^*M}$ is characterized by the relation

$$(4.2.1) \quad f(Q)_y u = Q_x^* u \quad \text{in } \mathcal{L},$$

where Q^* denotes the adjoint operator, and (\mathcal{L}, u) is a simple module along the conormal bundle to the diagonal Δ in $T^*(M \times M)$ (see [18]). Denote by \mathcal{C}_Δ the sheaf of complex microfunctions along the conormal bundle to Δ . By [18, Theorem 8.21], there exist $\lambda \in \mathbb{C}$ and an isomorphism

$$\varphi: \mathcal{E}_{\dot{T}^*M \times \dot{T}^*M}^{[\lambda]} \otimes_{\mathcal{E}_{\dot{T}^*M \times \dot{T}^*M}} \mathcal{C}_\Delta \xrightarrow{\sim} \mathcal{L},$$

so that $\varphi(P_y \otimes \delta_\Delta) = u$ for some invertible $P \in \mathcal{E}_{\dot{T}^*M}(\lambda)$. One then has

$$\begin{aligned} P_y Q_y P_y^{-1} u &= P_y Q_y P_y^{-1} \varphi(P_y \otimes \delta_\Delta) = \varphi(P_y Q_y \otimes \delta_\Delta) = \varphi(Q_x^* P_x^* \otimes \delta_\Delta) \\ &= Q_x^* \varphi(P_x^* \otimes \delta_\Delta) = Q_x^* \varphi(P_y \otimes \delta_\Delta) = Q_x^* u. \end{aligned}$$

It follows by (4.2.1) that one has $f = \text{Ad}(P)$. \square

4.3. Invertible \mathcal{E} -bimodules. Denote by P^*M the projective cotangent bundle of M and by $\gamma: \dot{T}^*M \rightarrow P^*M$ the projection. Set

$$\mathcal{E}_{P^*M} = \gamma_* \mathcal{E}_{\dot{T}^*M}.$$

This is a sheaf of \mathbb{C} -algebras endowed with a \mathbb{Z} -filtration such that $\mathcal{G}r \mathcal{E}_{P^*M} \simeq \bigoplus_{m \in \mathbb{Z}} \mathcal{O}_{P^*M}(m)$, where one sets $\mathcal{O}_{P^*M}(m) = \gamma_* \mathcal{O}_{\dot{T}^*M}(m)$. Note that $\mathcal{E}_{\dot{T}^*M}$ is constant along the fibers of γ . Since these are connected, the adjunction morphism gives an isomorphism

$$\gamma^{-1} \mathcal{E}_{P^*M} \xrightarrow{\sim} \mathcal{E}_{\dot{T}^*M}.$$

Lemma 4.3.1. *Let $Z \subset \dot{T}^*M$ be a closed conic analytic subset. Then*

$$H^j R\Gamma_Z \mathcal{E}_{\dot{T}^*M} = 0 \quad \text{for } j < \text{codim}_{\dot{T}^*M} Z.$$

Proof. Setting $W = \gamma(Z)$, we have $R\Gamma_Z \mathcal{E}_{\dot{T}^*M} \simeq \gamma^{-1} R\Gamma_W \mathcal{E}_{P^*M}$. We thus have to show that $H^j R\Gamma_W \mathcal{E}_{P^*M} = 0$ for $j < \text{codim}_{P^*M} W$. Identify \mathcal{E}_{P^*M} with the sheaf \mathcal{C}_Δ of complex microfunctions along the conormal bundle to the diagonal in $P^* = P^*(M \times M)$. By quantized contact transformations, \mathcal{C}_Δ can further be identified with the sheaf of complex microfunctions \mathcal{C}_S along the conormal bundle to a hypersurface $S \subset P^*$. One has $\mathcal{C}_S \simeq \mathcal{O}_S \oplus H_{[S]}^1 \mathcal{O}_{P^*} \simeq \mathcal{O}_S^{\oplus \mathbb{Z}}$. Hence $H^j R\Gamma_W \mathcal{C}_S = 0$ for $j < \text{codim}_S W$. \square

Proposition 4.3.2. *Let \mathcal{M} be a coherent torsion-free $\mathcal{E}_{\dot{T}^*M}$ -module. Then \mathcal{M} is locally free outside a closed conic analytic 2-codimensional subset.*

Proof. We will reduce to the analogous statement for \mathcal{O} -modules, which is well-known (see [25, Corollary 5.15]).

Set for short $\mathcal{E} = \mathcal{E}_{\dot{T}^*M}$, $\mathcal{E}(0) = \mathcal{E}_{\dot{T}^*M}(0)$ and $\mathcal{O}(0) = \mathcal{O}_{\dot{T}^*M}(0)$. A coherent $\mathcal{E}(0)$ -submodule $\mathcal{L} \subset \mathcal{M}$ such that $\mathcal{E}\mathcal{L} = \mathcal{M}$ is called a lattice.

(a) \mathcal{M} has a torsion-free lattice \mathcal{L} . In fact, let \mathcal{F} be a lattice in $\mathcal{M}^* = \text{Hom}_{\mathcal{E}}(\mathcal{M}, \mathcal{E})$. Then $\mathcal{F}^* = \text{Hom}_{\mathcal{E}(0)}(\mathcal{F}, \mathcal{E}(0)) \subset \text{Hom}_{\mathcal{E}}(\mathcal{M}^*, \mathcal{E}) = \mathcal{M}^{**}$ and $\mathcal{E}\mathcal{F}^* = \mathcal{M}^{**}$, i.e. \mathcal{F}^* is a lattice in \mathcal{M}^{**} . Then $\mathcal{L} = \mathcal{F}^* \cap \mathcal{M}$ is a lattice in \mathcal{M} . Since \mathcal{F}^* is reflexive (that is, $\mathcal{F}^* \rightarrow (\mathcal{F}^*)^{**}$ is an isomorphism), \mathcal{F}^* is torsion free, and so is its submodule \mathcal{L} .

(b) The coherent $\mathcal{O}(0)$ -module $\bar{\mathcal{L}} = \mathcal{L}/\mathcal{L}(-1)$ is torsion-free. In fact, consider the exact sequence

$$0 \rightarrow \mathcal{E}(-1) \rightarrow \mathcal{E}(0) \xrightarrow{\sigma_0} \mathcal{O}(0) \rightarrow 0.$$

Then $\mathcal{O}(0) \otimes_{\mathcal{E}(0)} \mathcal{L} \simeq \overline{\mathcal{L}}$. Hence $(\overline{\mathcal{L}})^* = \mathcal{H}om_{\mathcal{O}(0)}(\overline{\mathcal{L}}, \mathcal{O}(0)) \simeq \mathcal{H}om_{\mathcal{O}(0)}(\mathcal{O}(0) \otimes_{\mathcal{E}(0)} \mathcal{L}, \mathcal{O}(0)) \simeq \mathcal{H}om_{\mathcal{E}(0)}(\mathcal{L}, \mathcal{O}(0))$. The exact sequence

$$0 \rightarrow \mathcal{H}om_{\mathcal{E}(0)}(\mathcal{L}, \mathcal{E}(-1)) \rightarrow \mathcal{H}om_{\mathcal{E}(0)}(\mathcal{L}, \mathcal{E}(0)) \rightarrow \mathcal{H}om_{\mathcal{E}(0)}(\mathcal{L}, \mathcal{O}(0))$$

thus reads

$$0 \rightarrow \mathcal{L}^*(-1) \rightarrow \mathcal{L}^* \rightarrow (\overline{\mathcal{L}})^*.$$

Hence $\overline{\mathcal{L}}^* \subset (\overline{\mathcal{L}})^*$. Then $\overline{\mathcal{L}} \subset \overline{\mathcal{L}}^{**} \subset (\overline{\mathcal{L}}^*)^* \xrightarrow{\sim} (\overline{\mathcal{L}}^*)^{***}$, so that $\overline{\mathcal{L}}$ is torsion-free.

(c) Since $\overline{\mathcal{L}}$ is torsion-free, it is locally free outside a closed conic analytic 2-codimensional subset S . Hence the same holds true for \mathcal{L} by Nakayama lemma. Thus $\mathcal{M} = \mathcal{E}\mathcal{L}$ is also locally free outside S . \square

Remark 4.3.3. Since projective $\mathcal{E}_{\dot{T}^*M}$ -modules are torsion-free, it follows that $\mathcal{E}_{\dot{T}^*M}$ is (coherent) projective-free if $\dim M = 1$. This is no more true if $\dim M > 1$.

Set

$$\mathcal{E}_{\dot{T}^*M}^e = \mathcal{E}_{\dot{T}^*M}^{\text{op}} \otimes_{\mathbb{C}} \mathcal{E}_{\dot{T}^*M}.$$

Note that, for $[\lambda], [\mu] \in \mathbb{C}/\mathbb{Z}$ the morphism of $\mathcal{E}_{\dot{T}^*M}^e$ -modules

$$\mathcal{E}_{\dot{T}^*M}^{[\lambda]} \otimes_{\mathcal{E}_{\dot{T}^*M}} \mathcal{E}_{\dot{T}^*M}^{[\mu]} \rightarrow \mathcal{E}_{\dot{T}^*M}^{[\lambda+\mu]}, \quad P \otimes Q \mapsto PQ$$

is an isomorphism. In particular, $\mathcal{E}_{\dot{T}^*M}^{[\lambda]}$ is an invertible $\mathcal{E}_{\dot{T}^*M}^e$ -module. Moreover, if $P \in \mathcal{E}_{\dot{T}^*M}(\lambda)$ has non-vanishing symbol on $V \subset \dot{T}^*M$, there is an isomorphism of \mathcal{E}_V^e -modules (where we use notation (2.4.2))

$$(4.3.1) \quad \text{Ad}(P^{-1})\mathcal{E}_V \xrightarrow{\sim} \mathcal{E}_V^{[\lambda]}, \quad Q \mapsto PQ.$$

Lemma 4.3.4. For $[\lambda], [\mu] \in \mathbb{C}/\mathbb{Z}$, one has

$$\mathcal{H}om_{\mathcal{E}_{\dot{T}^*M}^e}(\mathcal{E}_{\dot{T}^*M}^{[\lambda]}, \mathcal{E}_{\dot{T}^*M}^{[\mu]}) = \begin{cases} \mathbb{C}_{\dot{T}^*M} & \text{for } [\lambda] = [\mu], \\ 0 & \text{otherwise.} \end{cases}$$

Proof. The problem is local and we take a system $(x) = (x_1, \dots, x_n)$ of local coordinates in $V \subset \dot{T}^*M$ such that ∂_1 is invertible in V . By (4.3.1)

$$\begin{aligned} \mathcal{H}om_{\mathcal{E}_V^e}(\mathcal{E}_V^{[\lambda]}, \mathcal{E}_V^{[\mu]}) &\simeq \mathcal{H}om_{\mathcal{E}_V^e}(\text{Ad}(\partial_1^{-\lambda})\mathcal{E}_V, \text{Ad}(\partial_1^{-\mu})\mathcal{E}_V) \\ &\simeq \{P \in \mathcal{E}_V : P\partial_1^{-\lambda}Q\partial_1^\lambda = \partial_1^{-\mu}Q\partial_1^\mu P, \forall Q \in \mathcal{E}_V\}. \end{aligned}$$

Assume that there exists $P \neq 0$ as above. Taking for Q the operators ∂_1 , x_i and ∂_i , respectively, we deduce that $[P, \partial_1] = [P, x_i] = [P, \partial_i] = 0$ for $i = 2, \dots, n$. It follows that P only depends on ∂_1 . Noting that $[\partial_1^\lambda, x_1] = \lambda\partial_1^{\lambda-1}$ and taking $Q = x_1$, we get

$$[x_1, P] = (\mu - \lambda)P\partial_1^{-1}.$$

Write $P = \sum_{j \leq m} c_j \partial_1^j$ with $c_i \in \mathbb{C}$ and $c_m \neq 0$. Then the above equality gives $m = \mu - \lambda$ and $c_j = 0$ for $j < m$. \square

The following result was communicated to us by Masaki Kashiwara (refer to [24] for related results).

Theorem 4.3.5. *Any invertible $\mathcal{E}_{\dot{T}^*M}^e$ -module is isomorphic to $L \otimes_{\mathbb{C}} \mathcal{E}_{\dot{T}^*M}^{[\lambda]}$, for some local system of rank one L and some locally constant \mathbb{C}/\mathbb{Z} -valued function $[\lambda]$.*

Proof. Set for short $\mathcal{E} = \mathcal{E}_{\dot{T}^*M}$. Let \mathcal{P} be an invertible \mathcal{E}^e -module. It is enough to show that \mathcal{P} is locally isomorphic to $\mathcal{E}^{[\lambda]}$ for some locally constant function $[\lambda]$. In fact, it will follow from Lemma 4.3.4 that $L = \text{Hom}_{\mathcal{E}^e}(\mathcal{E}^{[\lambda]}, \mathcal{P})$ is a local system of rank one and $L \otimes_{\mathbb{C}} \mathcal{E}^{[\lambda]} \xrightarrow{\sim} \mathcal{P}$.

(a) Since \mathcal{P} is invertible, the underlying \mathcal{E} -module $\bullet\mathcal{P}$ is projective locally of finite presentation by (iv) and (v) in Section 3.3, and hence coherent torsion-free. By Proposition 4.3.2, $\bullet\mathcal{P}$ is locally free outside a closed analytic 2-codimensional subset Z . As \mathcal{P} is invertible, its rank is one.

(b) Suppose that $\bullet\mathcal{P}$ is free of rank one. Then there exists $[\lambda]$ such that $\mathcal{P}^{[-\lambda]} = \mathcal{P} \otimes_{\mathcal{E}^e} \mathcal{E}^{[-\lambda]}$ admits a regular generator, i.e. a generator u of $\bullet\mathcal{P}^{[-\lambda]}$ such that $Pu = uP$ for any $P \in \mathcal{E}$. Indeed, let t be a generator of $\bullet\mathcal{P}$ and let $f: \mathcal{E} \xrightarrow{\sim} \mathcal{E}$, be the \mathbb{C} -algebra isomorphism as in (3.3.2): $f(P) = Q$ for Q such that $tP = Qt$. By Proposition 4.2.2, f is locally of the form $\text{Ad}(P)$ for some $\lambda \in \mathbb{C}$ and $P \in \mathcal{E}(\lambda)$ with never vanishing symbol. Then $u = tP^{-1}$ is a regular generator of $\mathcal{P}^{[-\lambda]}$.

Let V be a contractible open neighborhood of a point in Z . We are left to show that if $\bullet\mathcal{P}$ is locally free of rank one on $V \setminus Z$, then $\bullet\mathcal{P}^{[-\lambda]}$ has a regular generator on V . It will follow that $\mathcal{P}|_V \simeq \mathcal{E}_V^{[\lambda]}$.

(c) Since local regular generators u of $\mathcal{P}^{[-\lambda]}$ are unique up to multiplicative constants, $\mathbb{C}u \subset \mathcal{P}^{[-\lambda]}$ defines a local system of rank one on $V \setminus Z$. As $V \setminus Z$ is simply connected, such local system is constant. Thus $\mathcal{P}^{[-\lambda]}$ has a regular generator u on $V \setminus Z$.

Consider the distinguished triangle

$$R\Gamma_Z \mathcal{P}^{[-\lambda]} \rightarrow \mathcal{P}^{[-\lambda]} \rightarrow R\Gamma_{V \setminus Z} \mathcal{P}^{[-\lambda]} \xrightarrow{+1}$$

Since $\mathcal{P}^{[-\lambda]}$ is invertible, then $\bullet\mathcal{P}^{[-\lambda]}$ is flat by (iv) in Section 3.3, so that

$$R\Gamma_Z(V; \mathcal{P}^{[-\lambda]}) \simeq R\Gamma(V; R\Gamma_Z \mathcal{E} \otimes_{\mathcal{E}} \mathcal{P}^{[-\lambda]}).$$

By Lemma 4.3.1 one gets $H^j R\Gamma_Z(V; \mathcal{P}^{[-\lambda]}) = 0$ for $j = 0, 1$. It follows that $\Gamma(V; \mathcal{P}^{[-\lambda]}) \xrightarrow{\sim} \Gamma(V \setminus Z; \mathcal{P}^{[-\lambda]})$, hence the generator u of $\bullet\mathcal{P}^{[-\lambda]}$ on $V \setminus Z$ extends uniquely to V .

□

In particular, since any $\mathcal{E}_{\dot{T}^*M}^{[\lambda]}$ is a locally free right $\mathcal{E}_{\dot{T}^*M}$ -module of rank one by (4.3.1), it follows that the \mathbb{C} -algebra $\mathcal{E}_{\dot{T}^*M}$ is Picard good.

Recall that the projection $\gamma: \dot{T}^*M \rightarrow P^*M$ is a principal \mathbb{C}^\times -bundle.

Theorem 4.3.6. *The \mathbb{C} -algebra \mathcal{E}_{P^*M} is Picard good.*

Proof. Let us prove that any invertible $\mathcal{E}_{P^*M}^e$ -module \mathcal{P} is locally free of rank one as right \mathcal{E}_{P^*M} -module.

Since this is a local problem, we may restrict to a contractible open subset $U \subset P^*M$, so that $\gamma^{-1}(U) \simeq U \times \mathbb{C}^\times$. The $\mathcal{E}_{\gamma^{-1}(U)}^e$ -module $\gamma^{-1}\mathcal{P}$ being invertible, by Theorem 4.3.5 one gets

$$\mathcal{P} \xrightarrow{\sim} \gamma_* \gamma^{-1} \mathcal{P} \simeq \gamma_*(L \otimes_{\mathbb{C}} \mathcal{E}_{\gamma^{-1}(U)}^{[\lambda]})$$

for some $[\lambda] \in \mathbb{C}/\mathbb{Z}$ and some local system of rank one L on $\gamma^{-1}(U)$ with monodromy $e^{-2\pi i \lambda}$ on \mathbb{C}^\times .

By restricting to $U' \subset U$, we may assume that there exists an invertible operator D of order 1. This defines an isomorphism of right $\mathcal{E}_{U'}$ -modules

$$\mathcal{E}_{U'} \xrightarrow{\sim} \gamma_*(L \otimes_{\mathbb{C}} \mathcal{E}_{\gamma^{-1}(U')}^{[\lambda]}) \quad Q \mapsto D^\lambda Q.$$

□

Note that, given a local system of rank one L and $[\lambda] \in \mathbb{C}/\mathbb{Z}$, one has $\gamma_*(L \otimes_{\mathbb{C}} \mathcal{E}_{T^*M}^{[\lambda]}) \neq 0$ if and only if the monodromy of L along the fiber of γ is given by $e^{-2\pi i \lambda}$. In particular, $\gamma_* \mathcal{E}_{T^*M}^{[\lambda]} = 0$ for any $[\lambda] \neq 0$.

5. MICRODIFFERENTIAL ALGEBROIDS

Here we state and prove our results on classification of \mathcal{E} -algebroids on a contact manifold.

5.1. Contact manifolds. Let X be a complex manifold of odd dimension, say $2n - 1$. Denote by \mathcal{O}_X the sheaf of holomorphic functions and by Ω_X^1 the sheaf of holomorphic 1-forms. A structure of (complex) contact manifold on X is the assignment of a holomorphic principal \mathbb{C}^\times -bundle $\gamma: Y \rightarrow X$, called symplectification, and of a holomorphic one-form $\alpha \in \Gamma(Y; \Omega_Y^1)$, called contact form, such that $\omega = d\alpha$ is symplectic (i.e. ω^n vanishes nowhere) and $i_\theta \alpha = 0$, $L_\theta \alpha = \alpha$. Here, θ denotes the infinitesimal generator of the action of \mathbb{C}^\times on Y , i_θ the contraction and L_θ the Lie derivative. One may consider α as a global section of $\Omega_X^1 \otimes_{\mathcal{O}_X} \mathcal{O}_X(1)$, where $\mathcal{O}_X(1)$ denotes the dual of the sheaf of sections of the line bundle $\mathbb{C} \times_{\mathbb{C}^\times} Y$.

Let M be a complex manifold of dimension n . Then P^*M has a natural contact structure given by the Liouville one-form on \dot{T}^*M and by the projection $\gamma: \dot{T}^*M \rightarrow P^*M$. By Darboux theorem, P^*M is a local model for a contact manifold X , meaning that there are an open cover $\{U_i\}_{i \in I}$ of X and contact embeddings (i.e. embeddings preserving the contact forms) $j_i: U_i \hookrightarrow P^*M$ for any $i \in I$.

A fundamental result by [37] asserts that contact transformations (i.e. bi-holomorphisms preserving the contact forms) can be locally quantized.

This means the following. Let N be another complex manifold of dimension n , $U \subset P^*M$ and $V \subset P^*N$ open subsets and $\chi: U \rightarrow V$ a contact transformation. Then any $x \in U$ has an open neighborhood U' such that there is a \mathbb{C} -algebra isomorphism $\chi^{-1}(\mathcal{E}_{P^*N}|_{\chi(U')}) \xrightarrow{\sim} \mathcal{E}_{P^*M}|_{U'}$.

Definition 5.1.1. An \mathcal{E} -algebra on a contact manifold X is a sheaf \mathcal{A} of \mathbb{C} -algebras such that there are an open cover $\{U_i\}_{i \in I}$ of X , contact embeddings $j_i: U_i \hookrightarrow P^*M$ and \mathbb{C} -algebra isomorphisms $\mathcal{A}|_{U_i} \simeq j_i^{-1}\mathcal{E}_{P^*M}$ for any $i \in I$.

Given an \mathcal{E} -algebra \mathcal{A} , the \mathbb{C} -algebra $\gamma^{-1}\mathcal{A}$ on Y satisfies $\gamma^{-1}\mathcal{A}|_{\gamma^{-1}(U_i)} \simeq \tilde{j}_i^{-1}\mathcal{E}_{\tilde{T}^*M}$ for $\tilde{j}_i: \gamma^{-1}(U_i) \rightarrow \tilde{T}^*M$ a homogeneous symplectic transformation lifting $j_i: U_i \hookrightarrow P^*M$. Note that Proposition 4.2.2 implies that the invertible $\gamma^{-1}\mathcal{A}^e$ -module $(\gamma^{-1}\mathcal{A})^{[\lambda]}$ is well-defined for any $[\lambda] \in \mathbb{C}/\mathbb{Z}$.

To quantize X in the strict sense means to endow it with an \mathcal{E} -algebra (see [2, 34]). This might not be possible in general. However, as we now recall, Kashiwara [17] proved that X is endowed with a canonical \mathcal{E} -algebroid.

5.2. Microdifferential algebroids.

Definition 5.2.1. (i) An \mathcal{E} -algebroid on X is a \mathbb{C} -algebroid \mathbf{A} such that for every open subset $U \subset X$ and any object $\alpha \in \mathbf{A}(U)$, the \mathbb{C} -algebra $\mathcal{E}nd_{\mathbf{A}}(\alpha)$ is an \mathcal{E} -algebra on U .
(ii) A stack of twisted \mathcal{E} -modules on X is a \mathbb{C} -stack \mathbf{M} such that there are an open cover $\{U_i\}_{i \in I}$ of X , \mathcal{E} -algebras \mathcal{E}_i on U_i and equivalences $\mathbf{M}|_{U_i} \approx_{\mathbb{C}} \mathbf{Mod}(\mathcal{E}_i)$ for any $i \in I$.

Note that a \mathbb{C} -stack \mathbf{A} is an \mathcal{E} -algebroid if and only if there are an open cover $\{U_i\}_{i \in I}$ of X , \mathcal{E} -algebras \mathcal{E}_i on U_i and equivalences $\mathbf{A}|_{U_i} \approx_{\mathbb{C}} \mathcal{E}_i^+$ for any $i \in I$. In particular, $\mathbf{Mod}(\mathbf{A})$ is a stack of twisted \mathcal{E} -modules.

Kashiwara's construction of the canonical \mathcal{E} -algebroid on X was performed by patching data as explained in Appendix A.2 (see [10] for a more intrinsic construction). More precisely, in [17] he proved the existence of an open cover $\mathcal{U} = \{U_i\}_{i \in I}$ of X , of \mathcal{E} -algebras \mathcal{E}_i on U_i , of isomorphisms of \mathbb{C} -algebras $f_{ij}: \mathcal{E}_j \rightarrow \mathcal{E}_i$ on U_{ij} and of sections $a_{ijk} \in \Gamma(U_{ijk}; \mathcal{E}_i(0)^\times)$, satisfying the cocycle condition

$$(5.2.1) \quad \begin{cases} f_{ij}f_{jk} = \text{Ad}(a_{ijk})f_{ik}, \\ a_{ijk}a_{ikl} = f_{ij}(a_{jkl})a_{ijl}. \end{cases}$$

By Proposition A.2.1 (i), this implies

Theorem 5.2.2 ([17]). *Any complex contact manifold X is endowed with a canonical \mathcal{E} -algebroid \mathbf{E}_X .*

It follows that a \mathbb{C} -stack on X is an \mathcal{E} -algebroid (resp. a stack of twisted \mathcal{E} -modules) if and only if it is locally \mathbb{C} -equivalent to \mathbf{E}_X (resp. to

$\text{Mod}(\mathbf{E}_X)$). In particular, if $X = P^*M$ then \mathbf{E}_{P^*M} is \mathbb{C} -equivalent to \mathcal{E}_{P^*M} , and \mathcal{E} -algebroids are \mathbb{C} -twisted forms of \mathcal{E}_{P^*M} .

Recall that an algebroid is Picard good if and only if so are the algebras that locally represent it. Hence, by Theorem 4.3.6 one gets that any \mathcal{E} -algebroid, and in particular \mathbf{E}_X , is Picard good. From Proposition 3.4.3, we thus deduce the following

Theorem 5.2.3. (i) *Two \mathcal{E} -algebroids are \mathbb{C} -equivalent if and only if they are Morita equivalent.*
(ii) *Any stack of twisted \mathcal{E} -modules is \mathbb{C} -equivalent to the stack of modules over an \mathcal{E} -algebroid.*

To classify \mathcal{E} -algebroids, we thus need to compute the first cohomology with value in the stack of 2-groups $\text{Aut}_{\mathbb{C}}(\mathbf{E}_X) \approx \text{Inv}(\mathbf{E}_X^e)_{\text{op}}$, where we set $\mathbf{E}_X^e = \mathbf{E}_X^{\text{op}} \otimes_{\mathbb{C}} \mathbf{E}_X$.

5.3. Geometry of $\gamma: Y \rightarrow X$.

Lemma 5.3.1. *For M an abelian group, there is a distinguished triangle*

$$M_X \rightarrow R\gamma_*M_Y \rightarrow M_X[-1] \xrightarrow{+1}$$

Proof. As the complex $R\gamma_*M_Y$ is concentrated in degrees $[0, 1]$, by truncation it is enough to prove the isomorphisms

$$(5.3.1) \quad H^i R\gamma_*M_Y \simeq M_X, \quad \text{for } i = 0, 1.$$

For $i = 0$ it is induced by the adjunction morphism $M_X \rightarrow R\gamma_*M_Y$.

Set $SY = Y/\mathbb{R}_{>0}$ and consider γ as the composite of $p: Y \rightarrow SY$ and $q: SY \rightarrow X$, which are principal bundle for the groups $\mathbb{R}_{>0}$ and S^1 , respectively. Note that $Rp_*M_Y \simeq M_{SY}$, so that $R\gamma_*M_Y \simeq Rq_*M_{SY} \simeq Rq_!M_{SY}$. The infinitesimal generator θ of the action of \mathbb{C}^\times on Y induces a trivialization of the relative orientation sheaf $or_{SY/X}$. Hence $q^!M_X \simeq M_{SY}[1]$. Then the isomorphism (5.3.1) for $i = 1$ is induced by the adjunction morphism $Rq_!M_{SY} \simeq Rq_!q^!M_X[-1] \rightarrow M_X[-1]$. \square

Let $M = \mathbb{C}^\times$. The induced long exact cohomology sequence gives:

$$H^1(Y; \mathbb{C}^\times) \xrightarrow{\mu_1} H^0(X; \mathbb{C}^\times) \xrightarrow{\delta} H^2(X; \mathbb{C}^\times) \xrightarrow{\gamma^\#} H^2(Y; \mathbb{C}^\times) \xrightarrow{\mu_2} H^1(X; \mathbb{C}^\times).$$

We can represent elements of H^0 by locally constant \mathbb{C}^\times -valued functions, elements of H^1 by isomorphism classes of local systems of rank one, and elements of H^2 by \mathbb{C} -equivalence classes of invertible \mathbb{C} -algebroids (see Lemma 2.5.4). Let us describe the above sequence in these terms (see also [15, Chapitre V §§3.1, 3.2]), where we use the notation $[\cdot]$ both for isomorphism and \mathbb{C} -equivalence classes.

For L a local systems of rank one on Y , $\mu_1([L])$ is the locally constant function on X giving the monodromy of L along the fibers of γ .

Recall that \mathbb{C}_Y^+ denotes the stack of local systems of rank one on Y .

- Lemma 5.3.2.** (i) *There is a group isomorphism $\pi_0(\gamma_*\mathbb{C}_Y^+) \simeq \mathbb{C}_X^\times$, where the group structure on the left-hand side is induced by $\otimes_{\mathbb{C}}$.*
(ii) *If \mathbf{D} is a \mathbb{C} -stack on Y , then $\pi_0(\gamma_*\mathbf{D})$ is a \mathbb{C}_X^\times -sheaf (i.e., it is endowed with a \mathbb{C}_X^\times -action).*
(iii) *If \mathbf{T} is an invertible \mathbb{C}_Y -algebroid, then $\pi_0(\gamma_*\mathbf{T})$ is a \mathbb{C}_X^\times -torsor.*

Proof. (i) Recall that $\mathbb{C}_Y^\times[1]$ denotes the stack of \mathbb{C}_Y^\times -torsors. The functor

$$\mathbb{C}_Y^\times[1] \rightarrow \mathbb{C}_Y^+, \quad \mathcal{P} \mapsto \mathbb{C} \times_{\mathbb{C}^\times} \mathcal{P}$$

defines a group isomorphism $\pi_0(\gamma_*\mathbb{C}_Y^+) \simeq \pi_0(\gamma_*(\mathbb{C}_Y^\times[1]))$. By (1.4.3), the latter is isomorphic to $R^1\gamma_*\mathbb{C}_Y^\times$, hence to \mathbb{C}_X^\times by Lemma 5.3.1.

(ii) By using (2.1.1), one gets a \mathbb{C} -functor

$$\gamma_*\mathbb{C}_Y^+ \otimes_{\mathbb{C}} \gamma_*\mathbf{D} \rightarrow \gamma_*\mathbf{D}, \quad (L, \delta) \mapsto L \otimes_{\mathbb{C}} \delta.$$

This defines an action of $\pi_0(\gamma_*\mathbb{C}_Y^+) \simeq \mathbb{C}_X^\times$ on $\pi_0(\gamma_*\mathbf{D})$.

(iii) Since $R^2\gamma_*\mathbb{C}_Y^\times = 0$, the stack $\gamma_*\mathbf{T}$ is locally \mathbb{C} -equivalent to $\gamma_*\mathbb{C}_Y^+$. Hence $\pi_0(\gamma_*\mathbf{T})$ is a \mathbb{C}_X^\times -torsor by (i) and (ii). \square

Notation 5.3.3. Let \mathbf{C} be a \mathbb{C} -stack on X . For s a global section of $\pi_0(\mathbf{C})$, we denote by \mathbf{C}^s the full substack of \mathbf{C} whose objects c satisfy $[c] = s$ in $\pi_0(\mathbf{C})$.

Note that \mathbf{C}^s is a \mathbb{C} -algebroid, since $\pi_0(\mathbf{C}^s) = \{s\}_X$. It is locally \mathbb{C} -equivalent to the algebra $\mathcal{E}nd_{\mathbf{C}}(c)$ for any local representative c of s .

For $m \in H^0(X; \mathbb{C}^\times) \simeq \Gamma(X; \pi_0(\gamma_*\mathbb{C}_Y^+))$, one has

$$\delta(m) = [(\gamma_*\mathbb{C}_Y^+)^m].$$

Here, $(\gamma_*\mathbb{C}_Y^+)^m$ is identified with the \mathbb{C}_X -algebroid of local systems $L \in \gamma_*\mathbb{C}_Y^+$ with $\mu_1([L]) = m$. In particular, for $m = 1$ the \mathbb{C}_X -algebroid $(\gamma_*\mathbb{C}_Y^+)^1$ is equivalent to \mathbb{C}_X^+ via the adjunction functor $\mathbb{C}_X^+ \rightarrow \gamma_*\mathbb{C}_Y^+$. Moreover, one has a decomposition $\gamma_*\mathbb{C}_Y^+ \approx_{\mathbb{C}} \coprod_{m \in \mathbb{C}_X^\times} (\gamma_*\mathbb{C}_Y^+)^m$.

For \mathbf{S} an invertible \mathbb{C}_X -algebroid, $\gamma^\#([\mathbf{S}]) = [\gamma^{-1}\mathbf{S}]$.

Proposition 5.3.4. *For \mathbf{T} an invertible \mathbb{C}_Y -algebroid, $\mu_2([\mathbf{T}])$ is the class of the local system of rank one $\mathbb{C} \times_{\mathbb{C}^\times} \pi_0(\gamma_*\mathbf{T})$.*

Proof. By Lemma 5.3.2 (iii), $\pi_0(\gamma_*\mathbf{T})$ is a \mathbb{C}_X^\times -torsor. It follows that $\mathbb{C} \times_{\mathbb{C}^\times} \pi_0(\gamma_*\mathbf{T})$ is a local system of rank one on X .

Choose an open covering $\{U_i\}$ of X in such a way that \mathbf{T} is described, by means of Proposition A.1.1 (i), by the data $(\mathbb{C}_{V_i}^+, (\cdot) \otimes_{\mathbb{C}} M_{ji}, \mathbf{a}_{ijk})$, where $V_i = \gamma^{-1}(U_i)$ and M_{ji} are local systems of rank one on V_{ij} . Then $\mathbb{C} \times_{\mathbb{C}^\times} \pi_0(\gamma_*\mathbf{T})$ is represented by the 1-cocycle $\{\mu_1([M_{ji}])\}$ with values in \mathbb{C}^\times , which gives a Čech representative of the class $\mu_2([\mathbf{T}])$. \square

5.4. Classification results. Set

$$E_Y = \gamma^{-1}E_X, \quad E_Y^e = E_Y^{\text{op}} \otimes_{\mathbb{C}} E_Y \approx \gamma^{-1}(E_X^e).$$

Note that E_Y can be described by patching the \mathbb{C} -algebras $\gamma^{-1}\mathcal{E}_i$ along the pull back on Y of the data (5.2.1).

For $[\lambda] \in \mathbb{C}/\mathbb{Z}$, the algebroid version of the invertible bimodule $\mathcal{E}_{T^*M}^{[\lambda]}$ is the E_Y^e -module

$$E_Y^{[\lambda]} \in \text{Fct}_{\mathbb{C}_Y}(E_Y, E_Y) \subset \text{Mod}(E_Y^e)$$

locally defined by $(\cdot) \otimes_{\mathcal{E}_{T^*M}} \mathcal{E}_{T^*M}^{[\lambda]}$ (cp. Proposition 3.3.7).

Consider the direct image functor, obtained by using (1.1.1),

$$\gamma_*: \gamma_*\text{Mod}(E_Y^e) \rightarrow \text{Mod}(E_X^e)$$

and recall the morphism $H^1(Y; \mathbb{C}^\times) \xrightarrow{\mu_1} H^0(X; \mathbb{C}^\times) \simeq H^0(X; \mathbb{C}/\mathbb{Z})$ from §5.3.

Theorem 5.4.1. *The functor*

$$(5.4.1) \quad \gamma_*\text{Inv}(\mathbb{C}_Y) \rightarrow \text{Inv}(E_X^e), \quad L \mapsto \gamma_*(L \otimes_{\mathbb{C}} E_Y^{\mu_1(L^*)})$$

is an equivalence of stacks of 2-groups.

Proof. (a) A priori, $\gamma_*(L \otimes_{\mathbb{C}} E_Y^{\mu_1(L^*)})$ is an object of $\text{Mod}(E_X^e)$. It is locally, hence globally, invertible with inverse given by $\gamma_*(L^* \otimes_{\mathbb{C}} E_Y^{\mu_1(L)})$.

(b) The sheaf \mathbb{C}_Y is sent to E_X , since $\gamma_*(E_Y) \simeq E_X$ as E_X^e -modules. Moreover, the natural morphism

$$\gamma_*(L \otimes_{\mathbb{C}} E_Y^{\mu_1(L^*)}) \otimes_{E_X} \gamma_*(L' \otimes_{\mathbb{C}} E_Y^{\mu_1(L'^*)}) \rightarrow \gamma_*(L \otimes_{\mathbb{C}} L'^* \otimes_{\mathbb{C}} E_Y^{\mu_1(L^*) + \mu_1(L'^*)})$$

is locally, hence globally, an isomorphism. Hence (5.4.1) is monoidal.

(c) For an invertible E_X^e -module \mathcal{P} , define its exponent as the unique locally constant \mathbb{C}/\mathbb{Z} -valued function $\epsilon(\mathcal{P})$ on X such that $\gamma^{-1}\mathcal{P}$ is locally isomorphic to $E_Y^{\epsilon(\mathcal{P})}$ (this is well-defined by Theorem 4.3.5.). Then $\epsilon(\gamma_*(L \otimes_{\mathbb{C}} E_Y^{\mu_1(L^*)})) = \mu_1(L^*)$, and by using Lemma 4.3.4 one gets that the functor

$$\mathcal{P} \mapsto \text{Hom}_{E_Y^e}(E_Y^{\epsilon(\mathcal{P})}, \gamma^{-1}\mathcal{P})$$

is a quasi-inverse of (5.4.1). \square

Let $\text{Pic}(E_X)$ denote the set of isomorphism class of invertible E_X^e -modules, endowed with the group structure induced by \otimes_{E_X} .

Corollary 5.4.2. *There is a group isomorphism $\text{Pic}(E_X) \simeq H^1(Y; \mathbb{C}/\mathbb{Z})$.*

Theorem 5.4.3. *The set of \mathbb{C} -equivalence classes (resp. Morita classes) of \mathcal{E} -algebroids is canonically isomorphic, as a pointed set, to $H^2(Y; \mathbb{C}/\mathbb{Z})$.*

Proof. Since \mathbf{E}_X is Picard good, by Theorem 5.4.1 and Proposition 3.4.2 there is an equivalence of stacks of 2-groups

$$\mathrm{Aut}_{\mathbb{C}}(\mathbf{E}_X) \approx \gamma_* \mathrm{Inv}(\mathbb{C}_Y)_{\mathrm{op}}.$$

The right-hand term is equivalent to $\gamma_* \mathrm{Inv}(\mathbb{C}_Y)$ by the functor $L \mapsto L^*$. Since \mathbb{C}_Y is Picard good, from (3.4.3) and by using (1.4.2) one gets an equivalence of stacks of 2-groups

$$\gamma_* \mathrm{Inv}(\mathbb{C}_Y) \approx [R\gamma_* \mathbb{C}_Y^\times[1]].$$

It then follows from (1.4.1) that

$$(5.4.2) \quad H^1(X; \mathrm{Aut}_{\mathbb{C}}(\mathbf{E}_X)) \simeq H^2(Y; \mathbb{C}_Y^\times).$$

□

We end by giving a realization of isomorphism (5.4.2).

First, let us explain how to twist \mathbf{E}_Y by a local system of rank one L on X , obtaining a \mathbb{C} -algebroid \mathbf{E}_Y^L on Y locally \mathbb{C} -equivalent to \mathbf{E}_Y .

Choose an open covering $\{U_i\}$ of X in such a way that L is represented by a 1-cocycle $\{[\lambda_{ij}]\}$ with values in \mathbb{C}/\mathbb{Z} . Set $V_i = \gamma^{-1}(U_i)$ and consider the data $(\mathbf{E}_{V_i}, (\cdot) \otimes_{\mathbf{E}_{V_{ij}}} \mathbf{E}_{V_{ij}}^{[\lambda_{ij}]}, \mathbf{m}_{ijk})$, where \mathbf{m}_{ijk} denotes the invertible transformation induced by the canonical isomorphism of $\mathbf{E}_{V_{ijk}}^e$ -modules

$$\mathbf{E}_{V_{ijk}}^{[\lambda_{ij}]} \otimes_{\mathbf{E}_{V_{ijk}}} \mathbf{E}_{V_{ijk}}^{[\lambda_{jk}]} \xrightarrow{\sim} \mathbf{E}_{V_{ijk}}^{[\lambda_{ik}]}.$$

Then \mathbf{E}_Y^L is the \mathbb{C} -stack on Y obtained from these data by Proposition A.1.1 (i). Note that $(\mathbf{E}_Y^L)^{\mathrm{op}} \approx_{\mathbb{C}} \mathbf{E}_Y^{L^*}$ and $\mathbf{E}_Y^L \approx_{\mathbb{C}} \mathbf{E}_Y$ if L is trivial.

Denote by L^\times the \mathbb{C}_X^\times -torsor associated to L and recall from Lemma 5.3.2 that $\pi_0(\gamma_* \mathbf{E}_Y^L)$ is endowed with a \mathbb{C}_X^\times -action.

Lemma 5.4.4. $\pi_0(\gamma_* \mathbf{E}_Y^L) \simeq L^\times \times_{\mathbb{C}^\times} \pi_0(\gamma_* \mathbf{E}_Y)$ as \mathbb{C}_X^\times -sheaves.

Proof. Let $\{[\lambda_{ij}]\}$ be a 1-cocycle with values in \mathbb{C}/\mathbb{Z} representing L on an open covering $\{U_i\}$ of X . Then $\gamma_* \mathbf{E}_Y^L|_{U_i} \approx_{\mathbb{C}} \gamma_* \mathbf{E}_Y|_{U_i}$ and the associated glueing \mathbb{C} -equivalences $\gamma_* \mathbf{E}_Y|_{U_{ij}} \rightarrow \gamma_* \mathbf{E}_Y|_{U_{ij}}$ are given by $(\cdot) \otimes_{\mathbf{E}_{V_{ij}}} \mathbf{E}_{V_{ij}}^{[\lambda_{ij}]}$, where $V_i = \gamma^{-1}(U_i)$. We thus get isomorphisms of \mathbb{C}^\times -sheaves $\pi_0(\gamma_* \mathbf{E}_Y^L)|_{U_i} \simeq \pi_0(\gamma_* \mathbf{E}_Y)|_{U_i}$, with associated glueing automorphisms of $\pi_0(\gamma_* \mathbf{E}_Y)|_{U_{ij}}$ given by multiplication by $e^{2\pi i \lambda_{ij}}$. This follows from the commutative diagram of stacks of 2-groups

$$\begin{array}{ccc} \mathbb{C}/\mathbb{Z}_X[0] & \xrightarrow{\simeq} & \mathbb{C}_X^\times[0] \\ \downarrow & & \downarrow \\ \gamma_* \mathrm{Aut}_{\mathbb{C}}(\mathbf{E}_Y) & \xrightarrow{\pi_0} & \mathrm{Aut}(\pi_0(\gamma_* \mathbf{E}_Y))[0], \end{array}$$

where the left-hand vertical arrow is the functor $[\lambda] \mapsto (\cdot) \otimes_{E_Y} E_Y^{[\lambda]}$ and the right-hand one is the \mathbb{C}^\times -action. Hence $\pi_0(\gamma_* E_Y^L)$ is isomorphic to $\pi_0(\gamma_* E_Y)$ twisted by the \mathbb{C}_X^\times -torsor L^\times . \square

Let T be an invertible \mathbb{C}_Y -algebroid. Following Proposition 5.3.4, we denote by $\mu_2(T)$ the local system of rank one on X associated to the \mathbb{C}_X^\times -torsor $\pi_0(\gamma_* T)$.

Lemma 5.4.5. $\pi_0(\gamma_*(T \otimes_{\mathbb{C}} E_Y^{\mu_2(T^{op})})) \simeq \pi_0(\gamma_* E_Y)$ as \mathbb{C}_X^\times -sheaves.

Proof. By using the functor (2.1.1), one gets a morphism

$$\pi_0(\gamma_* T) \times \pi_0(\gamma_* E_Y^{\mu_2(T^{op})}) \rightarrow \pi_0(\gamma_*(T \otimes_{\mathbb{C}} E_Y^{\mu_2(T^{op})}))$$

which is \mathbb{C}^\times -equivariant on each term. Hence it factors through $\pi_0(\gamma_* T) \times_{\mathbb{C}^\times} \pi_0(\gamma_* E_Y^{\mu_2(T^{op})})$. By Lemma 5.4.5, this is isomorphic to $\pi_0(\gamma_* E_Y)$, since $\pi_0(\gamma_* T^{op})$ is isomorphic to the \mathbb{C}_X^\times -torsor opposite to $\pi_0(\gamma_* T)$. It follows that we have a morphism

$$\pi_0(\gamma_* E_Y) \rightarrow \pi_0(\gamma_*(T \otimes_{\mathbb{C}} E_Y^{\mu_2(T^{op})}))$$

of \mathbb{C}_X^\times -sheaves, which is locally, hence globally, an isomorphism. \square

Corollary 5.4.6. $\pi_0(\gamma_*(T \otimes_{\mathbb{C}} E_Y^{\mu_2(T^{op})}))$ has a canonical global section.

Proof. The adjunction functor $E_X \rightarrow \gamma_* E_Y$ defines a morphism $\pi_0(E_X) \rightarrow \pi_0(\gamma_* E_Y)$. Since $\pi_0(E_X)$ is the singleton-valued constant sheaf, this gives a global section of $\pi_0(\gamma_* E_Y)$, hence of $\pi_0(\gamma_*(T \otimes_{\mathbb{C}} E_Y^{\mu_2(T^{op})}))$ by Lemma 5.4.5. \square

Denote by *can* the canonical global section of $\pi_0(\gamma_*(T \otimes_{\mathbb{C}} E_Y^{\mu_2(T^{op})}))$. Then the inverse of the isomorphism (5.4.2) is realized as

$$[T] \mapsto [(\gamma_*(T \otimes_{\mathbb{C}} E_Y^{\mu_2(T^{op})}))^{can}],$$

where $[\cdot]$ denotes the \mathbb{C} -equivalence class and we use the Notation 5.3.3.

Assume that $T = \gamma^{-1}S$, for S an invertible \mathbb{C}_X -algebroid. Since $\mu_2(\gamma^{-1}S^{op})$ is trivial and $E_Y = \gamma^{-1}E_X$, the above isomorphism reduces to

$$[\gamma^{-1}S] \mapsto [(\gamma_* \gamma^{-1}(S \otimes_{\mathbb{C}} E_X))^{can}] = [S \otimes_{\mathbb{C}} E_X],$$

the latter being the class of the “twist” of E_X by S .

Remark 5.4.7. Replacing E_X by an \mathcal{E} -algebroid in the previous construction, one gets an action of $H^2(Y; \mathbb{C}_Y^\times)$ on the set of \mathbb{C} -equivalence classes (resp. Morita classes) of \mathcal{E} -algebroids. In such a way, the latter becomes an $H^2(Y; \mathbb{C}_Y^\times)$ -torsor and the canonical isomorphism (5.4.2) is obtained by choosing the \mathbb{C} -equivalence class of E_X as base point.

APPENDIX A. COCYCLES

For the reader's convenience, we recall here the descent conditions for stacks, and detail the case of algebroids. This is parallel to the case of gerbes. These results are well known and can be found for example in [15, 3, 17] (see also [26, 11, 35, 12, 4, 41]).

Let X be a topological space (or a site), and \mathcal{R} a sheaf of commutative rings on X . If $\mathcal{U} = \{U_i\}_{i \in I}$ is an open cover of X , we set $U_{ij} = U_i \cap U_j$, $U_{ijk} = U_i \cap U_j \cap U_k$ etc. We use the notation \bullet for the horizontal composition of transformations.

A.1. Glueing of stacks. Let us recall here how to recover \mathcal{R} -stacks, \mathcal{R} -functors and transformations from collections of local data.

Proposition A.1.1. *Let $\mathcal{U} = \{U_i\}_{i \in I}$ be an open cover of X .*

- (i) *Consider the descent datum $(C_i, f_{ij}, a_{ijk})_{ijk \in I}$, that is, C_i are \mathcal{R} -stacks on U_i , $f_{ij}: C_j \rightarrow C_i$ are \mathcal{R} -equivalences on U_{ij} and $a_{ijk}: f_{ik} \rightarrow f_{ij} \circ f_{jk}$ are invertible transformations on U_{ijk} , such that*

$$(A.1.1) \quad \begin{array}{ccc} f_{ij} \circ f_{jk} \circ f_{kl} & \xleftarrow{a_{ijk} \bullet \text{id}_{f_{kl}}} & f_{ik} \circ f_{kl} \\ \text{id}_{f_{ij}} \bullet a_{jkl} \uparrow & & \uparrow a_{ikl} \\ f_{ij} \circ f_{jl} & \xleftarrow{a_{ijl}} & f_{il} \end{array} \quad \text{commutes.}$$

Then there exists an \mathcal{R} -stack C on X endowed with \mathcal{R} -equivalences $f_i: C|_{U_i} \rightarrow C_i$ and invertible transformations $a_{ij}: f_i \rightarrow f_{ij} \circ f_j$ on U_{ij} , such that

$$\begin{array}{ccc} f_{ij} \circ f_{jk} \circ f_k & \xleftarrow{\text{id}_{f_{ij}} \bullet a_{jk}} & f_{ij} \circ f_j \\ a_{jkl} \bullet \text{id}_{f_k} \uparrow & & \uparrow a_{ij} \\ f_{ik} \circ f_k & \xleftarrow{a_{ik}} & f_i. \end{array} \quad \text{commutes.}$$

The \mathcal{R} -stack C is unique up to an \mathcal{R} -equivalence unique up to a unique invertible transformation.

- (ii) *Let C be as above, and let C' be associated with the descent datum $(C'_i, f'_{ij}, a'_{ijk})_{ijk \in I}$. Consider the descent datum $(g_i, b_{ij})_{ij \in I}$, that is, $g_i: C_i \rightarrow C'_i$ are \mathcal{R} -functors on U_i and $b_{ij}: f'_{ij} \circ g_j \rightarrow g_i \circ f_{ij}$ are invertible transformations on U_{ij} , such that*

$$(A.1.2) \quad \begin{array}{ccccc} g_i \circ f_{ij} \circ f_{jk} & \xleftarrow{\text{id}_{g_i} \bullet a_{ijk}} & g_i \circ f_{ik} & & \\ b_{ij} \bullet \text{id}_{f_{jk}} \uparrow & & & \nwarrow b_{ik} & \\ f'_{ij} \circ g_j \circ f_{jk} & \xleftarrow{\text{id}_{f'_{ij}} \bullet b_{jk}} & f'_{ij} \circ f'_{jk} \circ g_k & \xleftarrow{a'_{ijk} \bullet \text{id}_{g_k}} & f'_{ik} \circ g_k \end{array} \quad \text{commutes.}$$

Then there exists an \mathcal{R} -functor $g: \mathcal{C} \rightarrow \mathcal{C}'$ endowed with invertible transformations $b_i: f'_i \circ g \rightarrow g_i \circ f_i$ on U_i , such that

$$\begin{array}{ccccc}
 g_i \circ f_{ij} \circ f_j & \xleftarrow{\text{id}_{g_i} \bullet a_{ij}} & g_i \circ f_i & & \\
 \uparrow b_{ij} \bullet \text{id}_{f_j} & & \nwarrow b_i & & \\
 f'_{ij} \circ g_j \circ f_j & \xleftarrow{\text{id}_{f'_{ij}} \bullet b_j} & f'_{ij} \circ f'_j \circ g & \xleftarrow{a'_{ij} \bullet \text{id}_g} & f'_i \circ g
 \end{array}
 \quad \text{commutes.}$$

The \mathcal{R} -functor $g: \mathcal{C} \rightarrow \mathcal{C}'$ is unique up to a unique invertible transformation.

- (iii) Let $g: \mathcal{C} \rightarrow \mathcal{C}'$ be as above, and let $g': \mathcal{C} \rightarrow \mathcal{C}'$ be the \mathcal{R} -functor associated with the descent datum $(g'_i, b'_{ij})_{ij \in I}$. Consider the descent datum $(d_i)_{i \in I}$, that is, $d_i: g_i \rightarrow g'_i$ are transformations on U_i such that

$$\begin{array}{ccc}
 g_i \circ f_{ij} & \xleftarrow{b_{ij}} & f'_{ij} \circ g_j \\
 d_i \bullet \text{id}_{f_{ij}} \downarrow & & \downarrow \text{id}_{f'_{ij}} \bullet d_j \\
 g'_i \circ f_{ij} & \xleftarrow{b'_{ij}} & f'_{ij} \circ g'_j
 \end{array}
 \quad \text{commutes.}$$

(A.1.3)

Then, there exists a unique transformation $d: g \rightarrow g'$ such that $d|_{U_i} = d_i$.

Remark A.1.2. Let $\mathcal{V} = \{V_i\}_{i \in J}$ be open cover of X finer than \mathcal{U} , and choose a refinement map $\rho: J \rightarrow I$ (that is, $V_i \subset U_{\rho(i)}$ for any $i \in J$). Let $D = (C_i, f_{ij}, a_{ijk})_{ijk \in I}$ be a descent datum defined on \mathcal{U} and set

$$\tilde{C}_i = C_{\rho(i)}|_{V_i}, \quad \tilde{f}_{ij} = f_{\rho(i)\rho(j)}|_{V_{ij}}, \quad \tilde{a}_{ijk} = a_{\rho(i)\rho(j)\rho(k)}|_{V_{ijk}}.$$

Then $\rho^{-1}D = (\tilde{C}_i, \tilde{f}_{ij}, \tilde{a}_{ijk})_{ijk \in J}$ is a descent datum on \mathcal{V} which defines an \mathcal{R} -stack \mathcal{R} -equivalent to that associated to D .

Let $\rho': J \rightarrow I$ be another choice of a refinement map and $\rho'^{-1}D = (\tilde{C}'_i, \tilde{f}'_{ij}, \tilde{a}'_{ijk})_{ijk \in J}$ the associated descent datum on \mathcal{V} . Set

$$g_i = f_{\rho'(i)\rho(i)}|_{V_i}, \quad b_{ij} = (a_{\rho'(i)\rho'(j)\rho(i)}|_{V_{ij}} \bullet \text{id}_{\tilde{f}_{ij}})^{-1} \circ (\text{id}_{\tilde{f}'_{ij}} \bullet a_{\rho'(j)\rho(i)\rho(j)}|_{V_{ij}}).$$

Then $(g_i, b_{ij})_{ij \in J}$ is a descent datum as in (ii) and, since the g_i are equivalences, it defines an \mathcal{R} -equivalence between the \mathcal{R} -stacks associated to $\rho^{-1}D$ and $\rho'^{-1}D$.

A.2. Algebroid cocycles. We give here a description of \mathcal{R} -algebroids and \mathcal{R} -functors between them in terms of \mathcal{R} -algebras and \mathcal{R} -algebra morphisms.

Let A be an \mathcal{R} -algebroid on X . By definition, there exists an open cover $\{U_i\}_{i \in I}$ of X such that $A|_{U_i}$ is non-empty. For $\alpha_i \in A(U_i)$ and $\mathcal{A}_i = \text{End}_A(\alpha_i)$, there are \mathcal{R} -equivalences $f_i: A|_{U_i} \rightarrow \mathcal{A}_i^+$. Choose quasi-inverses f_i^{-1} and invertible transformations $\text{id} \rightarrow f_j^{-1} \circ f_j$. Set $f_{ij} := f_i \circ f_j^{-1}: \mathcal{A}_j^+ \rightarrow \mathcal{A}_i^+$ on U_{ij} . On U_{ijk} there are invertible transformations $a_{ijk}: f_{ik} \rightarrow f_{ij} \circ$

f_{jk} induced by $\text{id} \rightarrow f_j^{-1} \circ f_j$. On U_{ijkl} one checks that the diagram (A.1.1) commutes. By Proposition A.1.1 (i), the data $(\mathcal{A}_i, f_{ij}, a_{ijk})_{i,j,k \in I}$ are enough to reconstruct \mathbf{A} , in the sense that the stack obtained by glueing these data is \mathcal{R} -equivalent to \mathbf{A} .

The \mathcal{R} -equivalence $f_{ij}: \mathcal{A}_j^+ \rightarrow \mathcal{A}_i^+$ on U_{ij} is locally induced by \mathcal{R} -algebra isomorphisms. There thus exist an open cover $\{U_{ij}^\alpha\}_{\alpha \in A}$ of U_{ij} such that $f_{ij} = (f_{ij}^\alpha)^+$ on U_{ij}^α for $f_{ij}^\alpha: \mathcal{A}_j \rightarrow \mathcal{A}_i$ isomorphisms of \mathcal{R} -algebras. On triple intersections $U_{ijk}^{\alpha\beta\gamma} = U_{ij}^\alpha \cap U_{ik}^\beta \cap U_{jk}^\gamma$, the invertible transformations $a_{ijk}: (f_{ik}^\beta)^+ \rightarrow (f_{ij}^\alpha f_{jk}^\gamma)^+$ are given by invertible sections $a_{ijk}^{\alpha\beta\gamma} \in \mathcal{A}_i(U_{ijk}^{\alpha\beta\gamma})$ such that $f_{ij}^\alpha f_{jk}^\gamma = \text{Ad}(a_{ijk}^{\alpha\beta\gamma}) f_{ik}^\beta$. (Recall that we set $\text{Ad}(a)(b) = aba^{-1}$.) On quadruple intersections $U_{ijkl}^{\alpha\beta\gamma\delta\epsilon\varphi} = U_{ijk}^{\alpha\beta\gamma\delta\epsilon\varphi} \cap U_{ijl}^{\alpha\delta\epsilon} \cap U_{ikl}^{\beta\delta\epsilon\varphi} \cap U_{jkl}^{\gamma\epsilon\varphi}$, the commutative diagram (A.1.1) is equivalent to the equality $a_{ijk}^{\alpha\beta\gamma} a_{ikl}^{\beta\delta\epsilon\varphi} = f_{ij}^\alpha(a_{jkl}^{\gamma\epsilon\varphi}) a_{ijl}^{\alpha\delta\epsilon}$.

One can treat in the same manner \mathcal{R} -functors and transformations. We summarize the results in the next proposition. However, as indices of hypercovers are quite cumbersome, we will not write them explicitly anymore. Instead, we will assume that

(A.2.1) open covers of X are cofinal among hypercovers.

This is the case, for example, of paracompact spaces.

Proposition A.2.1. *Assume (A.2.1). Let $\{U_i\}_{i \in I}$ be a sufficiently fine open cover of X*

- (i) *Any \mathcal{R} -algebroid \mathbf{A} is reconstructed from a non-abelian cocycle $(\mathcal{A}_i, f_{ij}, a_{ijk})_{i,j,k \in I}$, that is, \mathcal{A}_i are \mathcal{R} -algebras on U_i , $f_{ij}: \mathcal{A}_j|_{U_{ij}} \rightarrow \mathcal{A}_i|_{U_{ij}}$ are \mathcal{R} -algebra isomorphisms and $a_{ijk} \in \mathcal{A}_i(U_{ijk})$ are invertible sections, such that*

$$(A.2.2) \quad \begin{cases} f_{ij} f_{jk} = \text{Ad}(a_{ijk}) f_{ik}, & \text{in } \mathcal{H}om_{\mathcal{R}\text{-Alg}_X}(\mathcal{A}_k, \mathcal{A}_i)(U_{ijk}), \\ a_{ijk} a_{ikl} = f_{ij}(a_{jkl}) a_{ijl}, & \text{in } \mathcal{A}_i(U_{ijkl}). \end{cases}$$

- (ii) *Let \mathbf{A} be as above, and let \mathbf{A}' be an \mathcal{R} -algebroid constructed from the non-abelian cocycle $(\mathcal{A}'_i, f'_{ij}, a'_{ijk})_{i,j,k \in I}$. Any \mathcal{R} -functor $\mathbf{g}: \mathbf{A} \rightarrow \mathbf{A}'$ is reconstructed from a non-abelian cocycle $(g_i, b_{ij})_{i,j \in I}$, that is, $g_i: \mathcal{A}_i \rightarrow \mathcal{A}'_i$ are \mathcal{R} -algebra morphisms and $b_{ij} \in \mathcal{A}'_i(U_{ij})$ are invertible sections, such that*

$$(A.2.3) \quad \begin{cases} g_i f_{ij} = \text{Ad}(b_{ij}) f'_{ij} g_j, & \text{in } \mathcal{H}om_{\mathcal{R}\text{-Alg}_X}(\mathcal{A}_j, \mathcal{A}'_i)(U_{ij}), \\ g_i(a_{ijk}) b_{ik} = b_{ij} f'_{ij}(b_{jk}) a'_{ijk}, & \text{in } \mathcal{A}'_i(U_{ijk}). \end{cases}$$

- (iii) *Let $\mathbf{g}: \mathbf{A} \rightarrow \mathbf{A}'$ be as above, and let $\mathbf{g}': \mathbf{A} \rightarrow \mathbf{A}'$ be constructed from the non-abelian cocycle $(g'_i, b'_{ij})_{i,j \in I}$. Any transformation of \mathcal{R} -functors $\mathbf{d}: \mathbf{g} \rightarrow \mathbf{g}'$ is reconstructed from a non-abelian cocycle*

$(d_i)_{i \in I}$, that is, $d_i \in \mathcal{A}'_i(U_i)$ are sections such that

$$(A.2.4) \quad d_i b_{ij} = b'_{ij} f'_{ij}(d_j), \quad \text{in } \mathcal{A}'_i(U_{ij}).$$

In particular, two non-abelian cocycles

$$(\mathcal{A}_i, f_{ij}, a_{ijk})_{i,j,k \in I}, \quad (\mathcal{A}'_i, f'_{ij}, a'_{ijk})_{i,j,k \in I},$$

are associated to \mathcal{R} -equivalent \mathcal{R} -algebroids if and only if, up to refinements, there exists a non-abelian cocycle $(g_i, b_{ij})_{i,j \in I}$ satisfying (A.2.3) with g_i isomorphisms of \mathcal{R} -algebras.

Viceversa, let \mathbf{A} be as in Proposition A.2.1 (i). For $i, j \in I$ let \mathcal{A}'_i be $\mathcal{R}|_{U_i}$ -algebras, $g_i: \mathcal{A}_i \rightarrow \mathcal{A}'_i$ isomorphisms of \mathcal{R} -algebras and $b_{ij} \in \mathcal{A}'_i(U_{ij})$ invertible sections. Then, the non-abelian cocycle $(\mathcal{A}'_i, f'_{ij}, a'_{ijk})_{i,j,k \in I}$ defined by (A.2.3) is associated to an \mathcal{R} -algebroid \mathcal{R} -equivalent to \mathbf{A} .

Remark A.2.2. Let $(\mathcal{A}_i, f_{ij}, a_{ijk})_{i,j,k \in I}$ be a non-abelian cocycle associated to an \mathcal{R} -algebroid \mathbf{A} . Then (A.2.2) implies the relations

$$f_{ii} = \text{Ad}(a_{iii}), \quad a_{iij} = a_{iii}, \quad a_{ijj} = f_{ij}(a_{jji}), \quad \text{for any } i, j \in I.$$

Setting $\mathcal{A}'_i = \mathcal{A}_i$, $g_i = \text{id}_{\mathcal{A}_i}$, and $b_{ij} = a_{iji}$ in (A.2.3), we get

$$f'_{ij} = \text{Ad}(a_{iji}^{-1})f_{ij}, \quad a'_{ijk} = f'_{ij}(a_{jki}^{-1}a_{jki}).$$

Then, the non-abelian cocycle $(\mathcal{A}'_i, f'_{ij}, a'_{ijk})_{i,j,k \in I}$ is associated to an \mathcal{R} -algebroid \mathcal{R} -equivalent to \mathbf{A} , and it satisfies the relations

$$f'_{ii} = \text{id}_{\mathcal{A}_i}, \quad a'_{iij} = a'_{ijj} = 1,$$

of a normalized cocycle in the sense of [3].

A.3. Module cocycles. Let \mathbf{A} be the \mathcal{R} -algebroid described over the open cover $\{U_i\}_{i \in I}$ of X by the non-abelian cocycle $(\mathcal{A}_i, f_{ij}, a_{ijk})_{i,j,k \in I}$. The stack of (left) \mathbf{A} -modules $\text{Mod}(\mathbf{A})$ is then described as in Proposition A.1.1 (i) by the family

$$(\text{Mod}(\mathcal{A}_i), \text{Mod}(f_{ji}^+), \text{Mod}(a_{kji}))_{i,j,k \in I}$$

(note the inversion of indices due to the fact that $\text{Mod}(\cdot)$ is contravariant). By Morita theory, the functor $\text{Mod}(f_{ji}^+) = f_{ji}(\cdot)$ is isomorphic to $\mathcal{P}_{ij} \otimes_{\mathcal{A}_j} (\cdot)$ for the invertible $\mathcal{A}_i \otimes_{\mathcal{R}} \mathcal{A}_j^{\text{op}}$ -module $\mathcal{P}_{ij} = f_{ji} \mathcal{A}_j$. We thus recover the description of twisted sheaves given in [19] (see also [17, 11, 13]).

Proposition A.3.1. *Let \mathbf{A} be as above. An object of $\text{Mod}(\mathbf{A})$ is described by a family $(\mathcal{M}_i, \varphi_{ij})_{i,j \in I}$, where $\mathcal{M}_i \in \text{Mod}(\mathcal{A}_i)$, and $\varphi_{ij} \in \text{Hom}_{\mathcal{A}_i}(f_{ji} \mathcal{M}_j|_{U_{ij}}, \mathcal{M}_i|_{U_{ij}})$ are isomorphisms, such that for any $u \in \mathcal{M}_k$ one has*

$$\varphi_{ij}(\varphi_{jk}(u)) = \varphi_{ik}(a_{kji}^{-1}u).$$

Proof. Let \mathcal{C} be an \mathcal{R} -stack as in Proposition A.1.1 (i). The statement follows by noticing that objects of $\mathcal{C}(X)$ are described by data

$$(\alpha_i, a_{ij})_{i,j \in I},$$

where $\alpha_i \in \mathcal{C}_i(U_i)$, and $a_{ij}: f_{ij}(\alpha_j) \rightarrow \alpha_i$ are isomorphisms in $\mathcal{C}_i(U_{ij})$, such that

$$a_{ij} \circ f_{ij}(a_{jk}) = a_{ik} \circ a_{ijk}^{-1}(\alpha_k)$$

as isomorphisms $f_{ij}f_{jk}(\alpha_k) \xrightarrow{\sim} \alpha_i$ in $\mathcal{C}_i(U_{ijk})$. \square

REFERENCES

- [1] E. Aldrovandi and B. Noohi, *Butterflies. I. Morphisms of 2-group stacks*, Adv. Math. **221**, no. 3 (2009), 687–773.
- [2] L. Boutet de Monvel, *Complex star algebras*, Math. Phys. Anal. Geom. **2**, no. 2 (1999), 113–139.
- [3] L. Breen, *On the classification of 2-gerbes and 2-stacks*, Astérisque **225**, 1994.
- [4] P. Bressler, A. Gorokhovsky, R. Nest and B. Tsygan, *Deformation quantization of gerbes*, Adv. Math. **214**, Issue 1 (2007), 230–266.
- [5] ———, *Deformations of algebroid stacks*, Adv. Math. **226**, no. 4 (2011), 3018–3087.
- [6] H. Bursztyn, *Semiclassical Geometry of Quantum Line Bundles and Morita Equivalence of Star Products*, Internat. Math. Res. Notices, **16** (2002), 821–846.
- [7] H. Bursztyn, V. Dolgushev and S. Waldmann, *Morita equivalence and characteristic classes of star products*, J. Reine Angew. Math. **662** (2012), 95–163.
- [8] H. Bursztyn and S. Waldmann, *Bimodule deformations, Picard groups and contravariant connections*, K-theory **31** (2004), 1–37.
- [9] D. Calaque and G. Halbout, *Weak quantization of Poisson structures*, Journal of Geometry and Physics **61** (2011), 1401–1414.
- [10] A. D'Agnolo and M. Kashiwara, *On quantization of complex symplectic manifolds*, Commun. Math. Phys., **308** no. 1 (2011), 81–113.
- [11] A. D'Agnolo and P. Polesello, *Stacks of twisted modules and integral transforms*, in: Geometric aspects of Dwork theory. Vol. I, 463–507, Walter de Gruyter, 2004.
- [12] ———, *Deformation quantization of complex involutive submanifolds*, in: Noncommutative geometry and physics (Yokohama, 2004), 127–137, World Scientific, 2005.
- [13] A. D'Agnolo and P. Schapira, *Quantization of complex Lagrangian submanifolds*, Adv. Math. **213**, no. 1 (2007), 358–379.
- [14] P. Deligne, *La formule de dualité globale*, in: Théorie des Topos et Cohomologie Etale des Schémas (SGA 4.3), 481–587, Lecture Notes in Mathematics **305**, Springer, 1973.
- [15] J. Giraud, *Cohomologie non abélienne*, Grundlehren der Math. Wiss. **179**, Springer, 1971.
- [16] M. Kashiwara, *Introduction to microlocal analysis*, Enseign. Math. **32**, no. 3-4 (1986), pp. 227–259.
- [17] ———, *Quantization of contact manifolds*, Publ. Res. Inst. Math. Sci. **32**, no. 1 (1996), 1–7.
- [18] ———, *D-modules and Microlocal Calculus*, Translations of Mathematical Monographs **217**, American Math. Soc. (2003).

- [19] ———, *Equivariant derived category and representation of real semisimple Lie groups*, in: Representation theory and complex analysis, 137–234, Lecture Notes in Math., **1931**, Springer, 2008.
- [20] M. Kashiwara and R. Rouquier, *Microlocalization of rational Cherednik algebras*, Duke Math. J. **144**, no. 3 (2008), 525–573.
- [21] M. Kashiwara and P. Schapira, *Ind-sheaves*, Astérisque **271** (2001).
- [22] ———, *Categories and sheaves*, Grundlehren der Math. Wiss. **332**, Springer, 2006.
- [23] ———, *Deformation quantization modules*, Astérisque **345** (2012), xi+147 pp.
- [24] M. Kashiwara and K. Vilonen, *Microdifferential systems and the codimension-three conjecture*, Ann. of Math. **180**, no. 2 (2014).
- [25] S. Kobayashi, *Differential geometry of complex vector bundles*, Publ. of the MSJ **15**, Princeton University Press; Iwanami Shoten, 1987.
- [26] M. Kontsevich, *Deformation quantization of algebraic varieties*, in: Euro-Conference Moshé Flato, Part III (Dijon, 2000), Lett. Math. Phys. **56**, no. 3 (2001), 271–294.
- [27] W. Lowen, *Algebroid prestacks and deformations of ringed spaces*, Trans. Amer. Math. Soc. **360**, no. 3 (2008), 1631–1660.
- [28] W. Lowen and M. Van den Bergh, *Deformation theory of abelian categories*, Trans. Amer. Math. Soc. **358**, no. 12 (2006), 5441–5483.
- [29] B. Mitchell, *Theory of categories*, Pure and Applied Mathematics **XVII**, Academic Press, 1965.
- [30] ———, *Rings with several objects*, Adv. Math. **8** (1972), 1–161.
- [31] ———, *Separable algebroids*, Mem. Amer. Math. Soc. **57** (1985), no. 333, iv+96.
- [32] P. Polesello, *Uniqueness of Kashiwara’s quantization of complex contact manifolds*, in: Geometric and topological methods for quantum field theory, 233–246, Contemp. Math. **434**, Amer. Math. Soc., (2007).
- [33] ———, *Classification of deformation quantization algebroids on complex symplectic manifolds*, Publ. Res. Inst. Math. Sci. **44**, no. 3 (2008), 725–748.
- [34] ———, *On quantizations of complex contact manifolds*, Adv. Math. **268** (2015), 129–158.
- [35] P. Polesello and P. Schapira, *Stacks of quantization-deformation modules on complex symplectic manifolds*, Int. Math. Res. Notices **2004:49** (2004), 2637–2664.
- [36] L. Prelli, *Proper stacks*, Publ. Res. Inst. Math. Sci. **43**, no. 3 (2007), 525–533.
- [37] M. Sato, T. Kawai, and M. Kashiwara, *Microfunctions and pseudo-differential equations*, in: Hyperfunctions and pseudo-differential equations (Katata 1971), 265–529, Lecture Notes in Math. **287**, Springer (1973).
- [38] P. Schapira, *Microdifferential systems in the complex domain*, Grundlehren der Math. Wiss. **269**, Springer, 1985.
- [39] I. Waschkes, *Microlocal Riemann-Hilbert correspondence*, Publ. Res. Inst. Math. Sci. **41**, no. 1 (2005), 37–72.
- [40] A. Yekutieli, *Derived equivalences between associative deformations*, J. Pure Appl. Algebra **214**, no. 8 (2010), 1469–1476.
- [41] ———, *Twisted deformation quantization of algebraic varieties*, Adv. Math. **268** (2015), 241–305.

DIPARTIMENTO DI MATEMATICA PURA ED APPLICATA, UNIVERSITÀ DI PADOVA,
VIA TRIESTE 63, 35121 PADOVA, ITALY

E-mail address: `dagnolo@math.unipd.it`

DIPARTIMENTO DI MATEMATICA PURA ED APPLICATA, UNIVERSITÀ DI PADOVA,
VIA TRIESTE 63, 35121 PADOVA, ITALY

E-mail address: `pietro@math.unipd.it`